

**From the holdings of the Department of Special Collections , Princeton University Library,
Princeton, NJ**

These documents are either in the public domain, or can only be used for educational and research purposes (Fair use) as per U.S. Copyright law (text below). By accessing this file, all users agree that their use falls in the public domain or within fair use as defined by the copyright law. We do not charge any permission or usage fees for the publication of images of material in our collections, including those provided by Princeton University Library, via our website, catalog, or directly from staff. For more information regarding copyright, credit, and citation please review the information on the Copyright, Credit and Citation Guidelines page of our website.

<https://library.princeton.edu/special-collections/policies/copyright-credit-and-citation-guidelines>. If you have any questions, including how to order higher-quality images from the collection, contact Special Collections Public Services via our Ask Us! Form. <https://library.princeton.edu/special-collections/ask-us>

Doctoral Dissertations, 2011-2026

Mathematics

2020

Extending fibrations of knot complements to ribbon disk complements

Miller, Maggie

Princeton, NJ : Princeton University

<http://arks.princeton.edu/ark:/88435/dsp01w6634654p>

U.S. Copyright law: The copyright law of the United States (Title 17, §108, United States Code) governs the making of photocopies or other reproductions of copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be used for any purpose other than private study, scholarship, or research. If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of fair use that user may be liable for copyright infringement. This institution reserves the right to refuse to accept a copying order if, in its judgment, fulfillment of the order would involve a violation of copyright law. <https://www.copyright.gov/title17/title17.pdf>

EXTENDING FIBRATIONS OF KNOT
COMPLEMENTS TO RIBBON DISK
COMPLEMENTS

MAGGIE HALL MILLER

A DISSERTATION
PRESENTED TO THE FACULTY
OF PRINCETON UNIVERSITY
IN CANDIDACY FOR THE DEGREE
OF DOCTOR OF PHILOSOPHY

RECOMMENDED FOR ACCEPTANCE
BY THE DEPARTMENT OF
MATHEMATICS
ADVISER: DAVID GABAI

JUNE 2020

© Copyright by Maggie Miller, 2020.

All rights reserved.

Abstract

We show that if K is a fibered ribbon knot in $S^3 = \partial B^4$ bounding ribbon disk D , then with a transversality condition the fibration on $S^3 \setminus \nu(K)$ extends to a fibration of $B^4 \setminus \nu(D)$. This partially answers a question of Casson and Gordon (“If D is a homotopy-ribbon disk bounded by a fibered knot, is D fibered?”). In particular, we show the fibration always extends when D has exactly two local minima, extending a result of Scharlemann for the unknot. More generally, we construct movies of singular fibrations on 4-manifolds and describe a sufficient property of a movie to imply the underlying 4-manifold is fibered over S^1 .

Acknowledgements

The work in this thesis was completed in November 2018. I entered graduate school intending to study 3-dimensional topology, but quickly transitioned to dimension 4.

Thanks to my advisor, Dave Gabai, for many long, helpful conversations about this project and several others. He made time to meet with me even when he was busy as chair of the department or with his own mathematics; I appreciate the time he spent with me as I developed as a graduate student.

Thanks also to Dave Gay, Jeff Meier, and Alex Zupan, who included me in the trisections group as a young graduate student. They helped me find other people who were doing interesting things and eventually develop my own mathematical interests.

Thanks to Cameron Gordon, who advised my undergraduate thesis and motivated me to apply to (and move to) Princeton. Also thanks to Danny Ruberman, who helped me with technical questions and was very friendly and supportive. I think they both generally contribute to a positive climate in the subfield of geometric topology.

Thanks to my co-authors: Paolo Aceto, Nickolas A. Castro, Irving Dai, Mark C. Hughes, Gabriel Islambouli, András Juhász, Seungwon Kim, Michael R. Klug, Peter Lambert-Cole, Jeffrey Meier, Allison N. Miller, JungHwan Park, András I. Stipsicz, Maggy Tomova, Ian Zemke, and Alexander Zupan. I am glad that I had the opportunity to work with them.

Thanks to my friends Antonio, Boya, Clayton, Daniel, Gabe, Hannah, Henry(?), Jason, Lena, Levent, Lucia, Mac, Mark, Nick, Patrick, PLC, Sarah, and Seungwon (some of whom are also my co-authors) for lots of interesting mathematical and non-mathematical conversations. (Special thanks to Clayton, who I talked to very often, and was probably my best friend during graduate school.)

Finally, thanks to Irving Dai for all of our time together through both of our graduate careers. I am happy that we will also be together as postdoctoral scholars.

I am supported by the NSF GRFP, Grant No. DGE-1656466.

To my parents, who always encouraged me.

Contents

Abstract	iii
Acknowledgements	iv
List of Tables	vii
List of Figures	viii
1 Introduction	1
2 Movies of singular fibrations on 4-manifolds with boundary	11
2.1 Basic definitions	11
2.2 Singularity charts: how to fiber a 4-manifold	16
3 Building blocks: simple movies of fibrations	25
3.1 Adding or cancelling singular points	26
3.2 Movies on Z^4 where $h : Z^4 \rightarrow I$ has one critical point	30
3.3 Positioning singularities in a fibration	36
3.4 Cancelling interior and boundary singularities	41
3.5 Movies on $B^3 \times I \setminus \nu(\text{saddle, minimum, or maximum disk})$	44
3.6 Composite movies	49
3.7 Proof of Lemma 2.4	58
4 Extending the fibration on the knot complement to a fibration of the disk complement	62

4.1	Proof of Theorem 1.8	62
4.2	Remark on the generalized cancellation movies	64
4.3	Understanding the fibers of $B^4 \setminus \nu(D)$	67
4.4	Proof of Theorem 1.10	69
5	Conclusion: explicitly finding the fiber of $B^4 \setminus \nu(D)$	71
6	More examples	75
7	More questions and final comments	81
	Bibliography	85

List of Tables

2.1	The type of a singularity in a movie of singular fibrations.	18
-----	--	----

List of Figures

1.1	A diagram of a ribbon disk D with boundary the square knot	5
1.2	A schematic of a ribbon disk in B^4	5
1.3	We perturb a band in one leaf to be transverse to a fibration.	7
2.1	Neighborhoods of singularities of singular fibrations.	12
2.2	A singular fibration $\mathcal{F}$ on a singular 3-manifold M^3	15
2.3	Smooth vs. singular fibers.	17
2.4	The type of a singularity in a movie of singular fibrations.	19
2.5	Possible configurations of two or three arcs meeting at an endpoint in a valid singularity chart.	21
2.6	An example of a valid singularity chart.	22
2.7	A movie of singular fibrations near a critical point of the height function.	24
3.1	A diagram of an interior stabilization movie.	27
3.2	Schema of interior stabilization movies.	27
3.3	As t decreases, we stabilize or destabilize $\mathcal{F}_t$ at a point in $\partial h^{-1}(t)$. . .	28
3.4	Valid singularity charts for each interior stabilization/destabilization.	29
3.5	Valid singularity charts for each boundary stabilization/destabilization.	29
3.6	The first boundary fusion movie.	30
3.7	The second boundary fusion movie.	31
3.8	The third boundary fusion movie.	32

3.9	The fourth boundary fusion movie.	33
3.10	The first boundary birth movie.	34
3.11	The second boundary birth movie.	34
3.12	Valid singularity charts for the boundary fusion, compression, birth, death and interior singularity movies.	36
3.13	Diagram of positioning a dot or half-dot.	38
3.14	Diagram of positioning a cone along an arc γ	39
3.15	Ensuring a positioned cone is type II.	40
3.16	Valid singularity charts for the positioning movies.	41
3.17	Schema of the cone, half-cone death and cone, half-dot death.	43
3.18	Valid singularity charts for the cone, half-cone death/birth and cone, half-dot death/birth.	44
3.19	Diagram of a saddle movie.	46
3.20	Diagram of a minimum movie.	48
3.21	Singularity charts for the saddle, minimum, and maximum movies.	49
3.22	Uncancellable cones and dots.	50
3.23	Some leaves in a band movie.	53
3.24	One leaf in a band movie.	54
3.25	Three leaves in a band movie.	54
3.26	A detailed view of a cone singularity at the end of a band movie.	55
3.27	Some leaves in a disk movie.	57
3.28	Closeup of a dot singularity introduced during a disk movie.	58
3.29	Singularity charts for cancellation movies.	59
3.30	An example valid singularity chart.	60
3.31	A more complicated example of a valid singularity chart.	61
4.1	Schematic of V_θ for increasing θ	66
4.2	Finding a cone that can cancel a dot.	67

4.3	All 1-handles in the 3-dimensional fiber of $B^4 \setminus \nu(D)$ are geometrically cancelled.	68
5.1	Valid singularity chart of the fibration constructed on a ribbon disk complement.	72
5.2	Fibering the complement of a ribbon disk bounded by 8_{20}	74
5.3	Fibering the complements of two distinct ribbon disks bounded by $11n_{74}$	74
6.1	Some 3-minimum ribbon disks satisfying the transversality condition of Theorem 1.8.	77
6.2	2-minimum disks of Owens–Swenton for all but one of the knots in Figure 6.1.	78
6.3	Ribbon 2-bridge knots admit ribbon disks with exactly two minima.	78
6.4	There are only 5 fibered, ribbon 2-bridge knots (Family 1).	80
6.5	There are only 5 fibered, ribbon 2-bridge knots (Families 2 and 3).	80

Chapter 1

Introduction

In dimension 3.5, low-dimensional topologists have studied knots in S^3 which bound smooth disks embedded in B^4 . We recall some basic definitions.

Definition 1.1. A knot K in a 3-manifold M is the oriented image of a smooth embedding $\phi : S^1 \rightarrow M^3$. We generally consider knots only up to the equivalence relation of smooth ambient isotopy.

Definition 1.2. A knot K in S^3 bounding a disk D smoothly embedded in B^4 is said to be *slice*.

If for some D , inclusion causes $\pi_1(S^3 \setminus K)$ to surject onto $\pi_1(B^4 \setminus D)$, then K is said to be *homotopy-ribbon*.

If for some D , the radial height function of B^4 is Morse on D with no local maxima, then K is said to be *ribbon*.

From definitions, we have the following order of inclusion.

$$\{\text{Ribbon knots}\} \subset \{\text{Homotopy-ribbon knots}\} \subset \{\text{Slice knots}\}.$$

Whether any of these inclusions are proper is a long-standing open question.

Slice-ribbon conjecture [Fox [8]]. Every slice knot in S^3 is ribbon.

One challenge in studying this problem is that it is difficult to show that a knot K is slice without implying that K is ribbon, and conversely that it is difficult to obstruct K from being ribbon without obstructing K from being slice. Casson and Gordon [2] found an obstruction to a *fibred* knot K being homotopy-ribbon.

Definition 1.3. Let K be a knot in S^3 . Then K is *fibred* if $S^3 \setminus \nu(K)$ is a bundle $\mathring{\Sigma}_g \times_\phi S^1 = \mathring{\Sigma}_g \times [0, 1] / [(x, 1) \sim (\phi(x), 0)]$, where $\mathring{\Sigma}_g$ is a Seifert surface for K and $\phi : \mathring{\Sigma}_g \rightarrow \mathring{\Sigma}_g$ is a surface automorphism fixing $\partial \mathring{\Sigma}_g$ pointwise.

The map ϕ is said to be the *monodromy* of K . This surface automorphism is well-defined up to conjugacy in the mapping class group of $\mathring{\Sigma}_g$. Note ϕ is an automorphism of a surface with boundary, but can be extended to a map $\hat{\phi}$ of the closed surface $\Sigma_g = \mathring{\Sigma}_g \cup D^2$. We call $\hat{\phi}$ the closed monodromy of K . We now state the general result of Casson and Gordon.

Theorem 5.1 of [2]. A fibred knot K in a homology 3-sphere M is homotopy-ribbon if and only if its closed monodromy extends over a handlebody. In particular, if $M = S^3$ and K is fibred and homotopy-ribbon, then there is a homotopy 4-ball V and slice disk $D \subset V$ so that $(S^3, K) = \partial(V, D)$ and $V \setminus \nu(D)$ is fibred by handlebodies.

(Note we did not define what it means for a knot in a homology 3-sphere to be homotopy-ribbon.)

From Casson and Gordon's proof of the above theorem, it is not obvious if the fibration on $S^3 \setminus \nu(K)$ should extend over $B^4 \setminus \nu(D)$ for a fixed disk D , or even whether V is diffeomorphic to B^4 .

Question 1.4. *Is $V \cong B^4$?*

Question 1.5. *Fix a homotopy-ribbon disk D in B^4 with boundary a fibred knot. Is $B^4 \setminus \nu(D)$ fibred over S^1 ? (i.e. "is D fibred?")*

If the answer to Question 1.5 is “yes”, then the 3-dimensional fibers are immediately known to be handlebodies by the following result of Larson and Meier.

Theorem 1.1 of [16]. Let D be a fibered slice disk in B^4 with fiber H . Then D is homotopy-ribbon if and only if H is a handlebody.

Remark 1.6. Larson and Meier use a different (and without the slice-ribbon conjecture, possibly more restrictive) definition of homotopy-ribbon, which we might call *strongly homotopy-ribbon*. A disk E is strongly homotopy-ribbon if $B^4 \setminus \nu(E)$ admits a handle decomposition without 3- or 4-handles. Their proof works for our stated definition of homotopy-ribbon as well (see chapter 7), yielding (disk fibered by handlebodies $\implies$ strongly homotopy-ribbon) and (disk fibered and homotopy-ribbon $\implies$ fibers are handlebodies). We conclude that for the class of fibered disks in B^4 , homotopy-ribbon and strongly homotopy-ribbon are equivalent conditions.

Remark 1.7. We remark for the general reader that although the sets of fibered knots and ribbon knots in S^3 are very special (and their intersection even more so), the set of prime fibered ribbon knots is infinite. For example, the three-stranded pretzel knots $P(\pm 2, n, -n)$ for any odd $|n| > 1$ are prime, fibered [10], and ribbon. In fact, these knots even admit a ribbon disk with only two local minima (the index-1 critical point “cuts” the ± 2 -twisted strand).

There are a total of 74 prime fibered ribbon knots of fewer than thirteen crossings [4].

In this paper, we prove the following relevant theorem.

Theorem 1.8. Let $K \subset S^3$ be a fibered knot bounding a ribbon disk $D \subset B^4$. View the index-1 critical points of D as fission bands attached to K . If the bands can be isotoped to be transverse to the fibration on $S^3 \setminus \nu(K)$, then the fibration extends to a fibration of handlebodies on $B^4 \setminus \nu(D)$.

To parse Theorem 1.8, we must understand how bands attached to a knot can describe a ribbon disk. Let K be a fibered knot in S^3 . Fix a ribbon disk $D \subset B^4$ with $\partial D = K$. View $B^4 = S^3 \times [0, 3]/(S^3 \times 0 \sim \text{pt})$, where $S^3 = \partial B^4 = S^3 \times 3$. We call the projection function $h : B^4 \rightarrow [0, 3]$ the height function on B^4 . A point $(x, t) \in S^3 \times [0, 3]/\sim$ is said to be at height t .

Isotope D so that

- $D \subset S^3 \times [1, 3]$.
- For $t_1 > t_2 > \dots > t_n \in (2, 3)$, $D \cap (S^3 \times t_i) =$ a link with a band attached (referred to as a ribbon band). We will always view these bands as “fission” bands, meaning resolving the band increases the number of components of the underlying link. In this setting, this means we view the band at height t_i as attached to a link equivalent to $D \cap (S^3 \times (t_i + \epsilon))$.
- For $s_1 > \dots > s_{n+1} \in (1, 2)$, $D \cap (S^3 \times s_i)$ is an $(n + 1 - i)$ -component unlink and a disjoint disk (referred to as a minimum disk).
- For all other $u \in (s_{n+1}, 3]$, $D \cap (S^3 \times u)$ is a nonsingular link. For all $u \in [0, s_{n+1})$, $D \cap (S^3 \times u) = \emptyset$.

In plain English, we isotope the disk D so that, starting from the boundary $S^3 = S^3 \times 3$ and moving downward, we first see all the saddle points and then the minima of D . We project the ribbon bands into $h^{-1}(0)$ to obtain a ribbon diagram $\mathcal{D}$ for D . (Assume generically that all ribbon bands are disjoint in $\mathcal{D}$.) Call the bands $b_1, \dots, b_n$, where b_i corresponds to the saddle at t_i .

See Figure 1.1 for an example of a ribbon disk and Figure 1.2 for a schematic of D in B^4 . Whenever we refer to a disk being determined by bands (or index-1 critical points of a disk corresponding to bands), we are referring to this description of a ribbon disk.

Now we discuss the transversality condition Theorem 1.8 places on fission bands.




Figure 1.1: Top: a diagram of a ribbon disk D with boundary the square knot. Below: Some cross-sections of $D \subset B^4$. Recall $h^{-1}(3)$ (that is, $t = 3$) is the boundary S^3 of B^4 .




Figure 1.2: A schematic of a ribbon disk in B^4 . In this example, the disk has 3 minima and 2 index-1 critical points (bands), as in Figure 1.1.

Definition 1.9. Let b be a band attached to a fibered knot K , where $S^3 \setminus \nu(K) = \mathring{\Sigma} \times I / \sim$. Say $b = \gamma \times [0, \epsilon]$ for some arc $\gamma \in S^3$ with $(\partial\gamma) \times [0, \epsilon] \subset K$. We say b is transverse to the fibration on $S^3 \setminus \nu(K)$ if $\mathring{\gamma}$ is transverse to every $\mathring{\Sigma} \times t$. Given a transverse band b , we may (and will) always assume $b \cap (\Sigma \times t) = \{\text{finite set of points}\} \times [0, \epsilon]$.

In Theorem 1.8, the hypothesis is that *all* bands in the description of D are disjoint and *simultaneously* transverse to the fibration on $S^3 \setminus \nu(K)$.

We will give an example and schematic of a ribbon disk in B^4 in chapter 4.

The following theorem follows almost immediately from the proof of Theorem 1.8.

Theorem 1.10. *Let K be a fibered knot in S^3 . Let $J \subset S^3$ be a knot so that there is a ribbon concordance from K to J (i.e. there is an annulus A properly embedded in $S^3 \times I$ with boundary $(K \times 1) \sqcup (\bar{J} \times 0)$ so that $\text{proj}_I|_A$ is Morse with no local maxima). Say the index-1 critical points of A correspond to fission bands attached to K . If the bands can be isotoped to be transverse to the fibration on $S^3 \setminus \nu(K)$, then $(S^3 \times I) \setminus \nu(A)$ is fibered by compression bodies.*

We remark that in Theorem 1.10, the fact that J is fibered follows immediately from [22, Prop. 5]. Theorem 1.10 is truly a 4-dimensional theorem about the ribbon concordance between K and J .

Both Theorems 1.8 and 1.10 are applications of our main theorem:

Theorem 1.11. *Let $\mathcal{F}_t$ be a valid movie of singular fibrations on Z^4 so that $\mathcal{F}_0$ and $\mathcal{F}_1$ are nonsingular. Then Z^4 is fibered over S^1 by $\{H_\theta\}$, where $H_\theta = \cup_t \mathcal{F}_t^{-1}(\theta)$.*

We will define a valid movie of singular fibrations in chapter 2.

Remark 1.12. The proof of Theorem 1.8 is constructive. From a minimum-genus surface $S \subset S^3$ for K and the fission bands defining D , it is possible to recover the attaching circles on S which define the handlebody to which the fibration extends. See chapter 5.




Figure 1.3: Top left: a band b lies in a single Seifert surface for a fibered knot K . Top right: Perturb the band near one component of $b \cap K$ so that $\mathring{b}$ is transverse to each leaf. (We can think of this perturbation as going halfway around $S^1 = \pi(S^3 \setminus K)$, where π is the projection map from $S^3 = \mathring{\Sigma} \times_{\phi} S^1$ to S^1 .) Bottom left: side view of b in one leaf, near one end of $b \cap K$. Bottom right: side view after perturbing b .

In Theorem 1.8, note that the fission bands are transverse to the open book on S^3 induced by the fibration of $S^3 \setminus \nu(K)$ in their *interior*. Near the binding of the open book on S^3 , the bands approach K asymptotically tangent to some leaf (see Fig. 1.3, bottom right).

Remark 1.13. If a fission band attached to a fibered knot K is contained in a fiber of $S^3 \setminus \nu(K)$, we may perturb the band to be transverse to the fibers as in Figure 1.3.

Now we discuss some corollaries of Theorem 1.8.

Corollary 1.14. *Let D be a ribbon disk in B^4 with exactly two minima. If ∂D is a fibered knot, then the fibration extends to a fibration of $B^4 \setminus \nu(D)$ by handlebodies.*

Proof. For a link J in S^3 , let $\chi(J) = \max\{\chi(S) \mid S \subset S^3 \text{ is an oriented surface bounded by } J \text{ with no closed components}\}$. Suppose J' is obtained from J by orientation-preserving band surgery along a band b . Then if $\chi(J') > \chi(J)$, the band b can be isotoped to lie in a Seifert surface S for J with $\chi(S) = \chi(J)$ [25, Prop 3.1].

Say $K = \partial D$. Let b be a fission band for K corresponding to the single saddle of D . Then surgering K along b yields the 2-component unlink $U \sqcup U$. Note $\chi(K) \leq$

$1 < 2 = \chi(U \sqcup U)$, so b may be isotoped to lie in a Seifert surface S for K with $\chi(S) = \chi(K)$. Then S is a fiber surface for K (see e.g. [1, Section 5B]), so the claim follows from Theorem 1.8 and Remark 1.13.

□

By gluing together two ribbon disks in separate 4-balls, we may obtain a 2-knot.

Definition 1.15. An n -knot is a copy of S^n smoothly embedded into S^{n+2} .

A 1-knot is a classical knot in S^3 , while a 2-knot is a knotted 2-sphere in S^4 .

Definition 1.16. Let D_1, D_2 be ribbon disks with $\partial D_1 = \partial D_2$. We say ∂D_i is an *equator* of the 2-knot $D_1 \cup \overline{D_2}$ in $S^4 = B^4 \cup \overline{B^4}$.

Note that an equator of a 2-knot G is not uniquely defined; for example, $K \# -K$ is an equator of the unknotted 2-sphere for any 1-knot K , as $K \# -K$ is naturally an equator of the 1-twist spin of K , which is unknotted [28].

If $G = D_1 \cup \overline{D_2}$ where D_1, D_2 have m_1, m_2 minima respectively, then in S^4 the 2-knot G can be taken to have m_1 minima, m_2 maxima, and $m_1 + m_2 - 2$ saddle points with respect to the standard height function. (Of course, ambient isotopy of S^4 may change the number of critical points induced by the standard height function on G .)

Corollary 1.17. *Let G be a 2-knot embedded in S^4 with exactly two minima, two maxima, and two saddle points with respect to the standard height function. If the equator of G (with respect to this decomposition) is a fibered 1-knot of genus- g , then G is fibered and the closure of the 3-dimensional fiber for G admits a genus- g Heegaard splitting.*

Proof. We have $G = D_1 \cup \overline{D_2}$, where each D_i has two minima and $K := \partial D_i$ is fibered. By Corollary 1.14, the fibration on $S^3 \setminus \nu(K)$ extends to fibrations on $B^4 \setminus \nu(D_1)$ and $B^4 \setminus \nu(D_2)$ by genus- g handlebodies. The claim follows. □

Corollary 1.14 can be viewed as an extension of the following fact due to Scharlemann [24]: If $D \subset B^4$ is a ribbon disk with two minima so that ∂D is the unknot, then D is boundary parallel (and hence $B^4 \setminus \nu(D) \cong S^1 \times B^3$ is fibered by 3-balls). Note that to prove Corollary 1.14, we used the fact that the saddle band lies in a minimum-genus surface of the equator. This fact also proves Scharlemann’s theorem, so we have extended this result but do not claim a simpler proof.

Finally, we remark that there is no analogue of Theorem 1.8 for links, as a fibered $(r > 1)$ -component link cannot be slice (in the sense of bounding r disjoint slice disks in B^4). This follows from the fact that for $r > 1$, a slice r -component link L has Alexander polynomial¹ $\Delta(t_1, \dots, t_r) = 0$ [7]. Meanwhile, if L is fibered, then $\Delta(t, \dots, t)$ must be monic, as $t^k(t-1)\Delta(t, \dots, t)$ is the characteristic polynomial of the monodromy of L for some $k \in \mathbb{Z}$ [21, Lemma 10.1].

More fundamentally, this result cannot be extended to links because for $r > 1$, $\chi(B^4 \setminus \nu(\sqcup_r D^2)) < 0$, so the complement of multiple ribbon disks in B^4 cannot be fibered.² Thus, Theorem 1.8 passes a basic sanity check for links.

Outline

- Chapter 2: We give basic definitions and prove Theorem 1.11.
- Chapter 3: We construct a library of basic movies of singular fibrations used to construct fibrations on larger 4-manifolds.
- Chapter 4: We prove Theorems 1.8 and 1.10. We mainly focus on Theorem 1.8, and then quickly prove Theorem 1.10 using the same argument.

¹We are referring to Fox’s definition of the Alexander polynomial of a link, as in [7].

²More generally, a multi-component link of S^k s embedded in S^{k+2} is *never* fibered for $k > 1$ [11]. For k even, this follows easily from the Euler characteristic argument.

- Chapter 5: We discuss how to find the 2-handle attaching circles on a fiber in $S^3 \setminus \nu(K)$ which define the handlebody fiber in $B^4 \setminus \nu(D)$ built during the construction of Theorem 1.8.
- Chapter 6: We provide examples of fibered ribbon disks. In particular, we produce a fibered ribbon disk for each prime fibered slice knot of fewer than 13 crossings.
- Chapter 7: We discuss relevant open questions.

Chapter 2

Movies of singular fibrations on 4-manifolds with boundary

2.1 Basic definitions

In this section, we essentially give alternate definitions of circular Morse functions and their critical points specifically in dimension three. These descriptions will be useful in explicit constructions.

Definition 2.1. A *singular fibration* $\mathcal{F}$ on a compact 3-manifold M^3 is a smooth map $\mathcal{F} : M^3 \rightarrow S^1$ with the following properties:

- Each $\mathcal{F}^{-1}(\theta)$ is a compact, properly embedded surface away from a finite number of *cone* or *dot* singularities in its interior and *half-cones*, *half-dots*, and *bowls* in its boundary (see below).
- For all but finitely many θ , $\mathcal{F}^{-1}(\theta)$ is a nonsingular surface.
- There are finitely many singularities in the foliation $\mathcal{F}$ induces on ∂M^3 . If $\mathcal{F}^{-1}(\theta) \cap \partial M$ contains such a singular point, then near that point $\mathcal{F}^{-1}(\theta)$ is a half-cone, a half-dot, or a bowl. See Figure 2.1.




Figure 2.1: Top: neighborhoods of a cone and dot singularity in $\mathring{M}^3$. Bottom: Leaves of $\mathcal{F}$ near a half-cone, half-dot, and bowl singular point in the foliation induced on ∂M^3 .

See Figure 2.1 for an illustration of cone and dot singularities, as well as singularities in the induced foliation on ∂M^3 . We believe this figure and the names of the singularities to be sufficient to understand what they are, but we include the following more precise descriptions for completeness. (These descriptions can be safely ignored if the reader is comfortable with Figure 2.1).

For $p \in \mathring{M}^3$, We say that $p \in \mathcal{F}^{-1}(\theta)$ is a *cone* singularity if for some small 3-ball neighborhood of p in M^3 , with coordinates x, y, z , we have $\mathcal{F}^{-1}(\theta) = \{x^2 + y^2 = z^2\}$, and for $\theta' \neq \theta$, $\mathcal{F}^{-1}(\theta') = \{x^2 + y^2 = z^2 + c_{\theta'}\}$ for some constant $c_{\theta'} \neq 0$. In words, near p the set $\mathcal{F}^{-1}(\theta)$ looks like a double cone, while each nearby $\mathcal{F}^{-1}(\theta')$ looks like a hyperboloid of one or two sheets.

For $p \in \mathring{M}^3$, we say $p \in \mathcal{F}^{-1}(\theta)$ is a *dot* singularity if for some small 3-ball neighborhood of p in M^3 , with coordinates x, y, z , we have $\mathcal{F}^{-1}(\theta) = \{(0, 0, 0)\}$ and for $\theta' \neq \theta$, $\mathcal{F}^{-1}(\theta') = \{x^2 + y^2 + z^2 = c_{\theta'}\}$ for some constant $c_{\theta'} \neq 0$. In words, near p the set $\mathcal{F}^{-1}(\theta)$ looks like an isolated point, while nearby $\mathcal{F}^{-1}(\theta')$ are nested spherical shells (or the empty set).

For $p \in \partial M^3$, we say $p \in \mathcal{F}^{-1}(\theta)$ is a *half-cone* singularity if for some small neighborhood of p in M^3 , with coordinates x, y, z , we have $\mathcal{F}^{-1}(\theta) = \{x^2 + z^2 =$

$y^2, z \geq 0\}$, and for $\theta' \neq \theta$, $\mathcal{F}^{-1}(\theta') = \{x^2 + z^2 = y^2 + c_{\theta'}, z \geq 0\}$ for some constant $c_{\theta'} \neq 0$. In words, near p the set $\mathcal{F}^{-1}(\theta)$ looks like half of a cone (half of each side), while each nearby $\mathcal{F}^{-1}(\theta')$ looks like half of a hyperboloid of one or two sheets.

For $p \in \partial M^3$, we say $p \in \mathcal{F}^{-1}(\theta)$ is a *half-dot* singularity if for some small neighborhood of p in M^3 , with coordinates x, y, z , we have $\mathcal{F}^{-1}(\theta) = \{(0, 0, 0)\}$ and for $\theta' \neq \theta$, $\mathcal{F}^{-1}(\theta') = \{x^2 + y^2 + z^2 = c_{\theta'}, z \geq 0\}$ for some constant $c_{\theta'} \neq 0$. In words, near p the set $\mathcal{F}^{-1}(\theta)$ looks like an isolated point, while nearby $\mathcal{F}^{-1}(\theta')$ are nested half-spherical shells (or the empty set).

For $p \in \partial M^3$, we say $p \in \mathcal{F}^{-1}(\theta)$ is a *bowl* singularity if for some small neighborhood of p in M^3 , with coordinates x, y, z , we have $\mathcal{F}^{-1}(\theta) = \{x^2 + y^2 = z\}$ and for $\theta' \neq \theta$, $\mathcal{F}^{-1}(\theta') = \{x^2 + y^2 = z + c_{\theta'}, z \geq 0\}$ for some constant $c_{\theta'} \neq 0$. In words, near p the set $\mathcal{F}^{-1}(\theta)$ looks like a bowl (i.e. a circular paraboloid), while nearby $\mathcal{F}^{-1}(\theta')$ are nested bowls or cylinders.

We may think of $\mathcal{F}$ as being a circle-valued Morse function on M^3 , so that the cone and dot singularities correspond to critical points in the Morse function $\mathcal{F}$. A cone corresponds to a critical point of index 1 or 2, while a dot to a critical point of index 0 or 3 (depending on the orientation of each $\mathcal{F}^{-1}(\theta)$ induced by the positive orientation of S^1).

We will refer to each $\mathcal{F}^{-1}(\theta)$ as a *leaf* of $\mathcal{F}$.

We may extend the definition of singular fibrations to some singular 3-manifolds.

Definition 2.2. Let M^3 be a closed, compact, singular 3-manifold with finitely many singularities $S = \{p_1, \dots, p_n, q_1, \dots, q_m \mid p_i \in \partial M^3, q_j \in \mathring{M}^3\}$. For $p_i \in S \cap \partial M^3$, we require p_i to have a neighborhood homeomorphic to one of: $\{x^2 + y^2 \geq z^2\}$, $\{x^2 + y^2 \leq z^2\}$, $\{x^2 + y^2 + z^2 \geq 0 \mid (0, 0, 0) \text{ is artificially declared to be a boundary point}\}$ or $\{(0, 0, 0)\}$.

For $q_i \in S \cap \mathring{M}^3$, we require q_i to have a neighborhood of the form $P \cup_{\nu(q_i) \cap \partial(P)} \overline{P}$, where P is a singular manifold with boundary and q_i is a singularity as above. (That

is, we first understand a neighborhood P of a singularity in ∂M^3 . Then an interior singularity has a neighborhood which is a double of P .)

This means ∂M^3 is the disjoint union of surfaces with a finite number of cone singularities and isolated points $\{p_1, \dots, p_n\}$.

Let $\mathcal{G}$ be a map $\mathcal{G} : (M^3 \setminus S) \rightarrow S^1$ which is a singular fibration on M^3 away from singularities of M^3 .

We extend $\mathcal{G}$ to $\mathcal{F} : M^3 \rightarrow S^1$ by saying $\mathcal{F}(x) = \theta$ if there is a smooth path $\gamma : [0, 1] \rightarrow M^3$ with $\gamma(1) = x$ and $\lim_{t \rightarrow 1} \mathcal{G}(\gamma(t)) = \theta$. If x is isolated in M^3 , we choose $\mathcal{F}(x)$ arbitrarily. We require that $\mathcal{G}$ be defined so that near a singular point p_i in ∂M^3 , $\mathcal{F}$ is as in one of the diagrams in Figure 2.2. Near a singular point $q_i \in \overset{\circ}{M}^3$, $\mathcal{F}$ must be as in two mirror copies of one diagram of Figure 2.2 glued along the boundary near q_i . In particular, we require $\mathcal{F}$ be a well-defined function.

We call $\mathcal{F}$ a singular fibration of M^3 .

Definition 2.3. A *movie of singular fibrations* on X^4 with Morse function $h : X^4 \rightarrow I$ is a family of smooth maps $\mathcal{F}_t : h^{-1}(t) \rightarrow S^1$ so that each $\mathcal{F}_t$ is a singular fibration on $h^{-1}(t)$ and the $\mathcal{F}_t$ vary smoothly with t (i.e. $x \mapsto \mathcal{F}_{h(x)}(x)$ is a smooth map from X^4 to S^1).

Lemma 2.4. *Let Z^4 be a compact 4-manifold. Fix a Morse function $h : Z^4 \rightarrow I$. Suppose $\mathcal{F}_t$ is a movie of singular fibrations on Z^4 (i.e. a family of smooth maps $\mathcal{F}_t : h^{-1}(t) \rightarrow S^1$) so that $\mathcal{F}_1$ and $\mathcal{F}_0$ are fibrations (with no singularities) and $\mathcal{F}_t$ is a concatenation of band, disk, and simple or generalized cancellation movies (to be defined in chapter 3) in order (decreasing t).*

Then $\mathcal{F}_1 \sqcup \mathcal{F}_0$ extends to a fibration on Z^4 .

We delay the proof of Lemma 2.4 until Section 3.7.

Our strategy for proving Theorem 1.8 is to construct a movie of singular fibrations on $B^4 \setminus \nu(D)$ from S^3 to just above the lowest minimum of D . We choose this movie




Figure 2.2: A singular fibration $\mathcal{F}$ on a singular 3-manifold M^3 , as in Def. 2.2. We require $\mathcal{F}$ agree with one of these six configurations in a neighborhood of a singular point in ∂M^3 . Near a singular point in $\overset{\circ}{M}^3$, $\mathcal{F}$ should agree with two copies of one of these diagrams, glued together along boundary near the singularity. **We first describe each figure as if the singularity is in ∂M^3 .** First four: the gray cone is contained in ∂M^3 . At the cone point, a leaf of $\mathcal{F}$ is either a wedge of two disks along a boundary point, a disk centered at the cone, a wedge of two disks along a boundary point, or a point (respectively). A nearby leaf is a disk, annulus, two disjoint disks, or disk (respectively). Last two: the black dot is an isolated point of ∂M^3 . We draw leaves of $\mathcal{F}$ near this point, in the case that the point is not isolated in M^3 , and then finally the case that the point is isolated in M^3 . **We now describe each figure as if the singularity is in $\overset{\circ}{M}^3$.** First four: at the singularity, a leaf of $\mathcal{F}$ is either a cone, a cone, a cone, or a point (respectively). A nearby leaf is a cylinder, cylinder, two disjoint disks, or sphere (respectively). Last two: the black dot is an isolated point of $\overset{\circ}{M}^3$. The leaf of $\mathcal{F}$ containing the singularity is either a cone or a point, respectively.

so that the top singular fibration on $S^3 \setminus \nu(K)$ is the honest fibration of the fibered knot K , and that the bottom singular fibration is the fibration of a solid torus by meridian disks. In the proof of Lemma 2.4, we will show this gives a fibration on $B^4 \setminus \nu(D)$. (Note that a movie of singular fibrations on $B^4 \setminus \nu(D)$ is *not* equivalent to a fibration of $B^4 \setminus \nu(D)$ – this is an additional step.) By [16], this immediately completes the proof, but we will separately analyze the constructed movie to directly show that the fibers of $B^4 \setminus \nu(D)$ are handlebodies.

To construct the movie, we give several basic building block movies of singular fibrations in chapter 3. First, we discuss how movies of singular fibrations may describe smooth fibrations of a 4-manifold.

2.2 Singularity charts: how to fiber a 4-manifold

Suppose we have a movie $\mathcal{F}_t$ of singular fibrations on Z^4 and we hope to find a fibration $\{H_\theta\}_{\theta \in S^1}$ of Z^4 where each $H_\theta = \cup_t \mathcal{F}_t^{-1}(g_t(\theta))$ for some $g_t : S^1 \rightarrow S^1$.

First, we consider two natural (and failing) attempts to create a fibration on Z^4 .

- The “vertical fiber” strategy: Let $H_\theta = \cup_t \mathcal{F}_t^{-1}(\theta)$. This fails because $\mathcal{F}_t^{-1}(\theta)$ may be singular for an interval of t values. Then the given union will yield a singular 3-manifold.
- The “tilted fiber” strategy: Let $H_\theta = \cup_t \mathcal{F}_t^{-1}(\theta + ct)$ for a constant c . For some choice of c , singular cross-sections of H_θ are isolated. However this strategy fails (in part) because we may change the direction in which the 3-dimensional fibers flow through a singular leaf in $\mathcal{F}_t$, leading to some 3-dimensional fiber being singular.

It is very important that the reader unstand the failing of the tilted fiber strategy. Attempt to take $H_\theta : \cup_t \mathcal{F}_t^{-1}(\theta + ct)$. Suppose p is a cone point of $\mathcal{F}_t$ for $t \in [s', s'']$.




Figure 2.3: Left and right: cross-sections of a smooth 3-manifold. Each has one singular cross-section (containing a cone). The nearby cross-sections are *different* resolutions of that cone. Middle: cross-sections of a *singular* 3-manifold. Even though there is only an isolated singular cross-section (a single cone), the resolutions of the cone are *the same* above and below the singular cross-section.

Suppose that $H_{\theta'}$ and $H_{\theta''}$ meet p at heights s' and s'' respectively. Suppose moreover that near p , $H_{\theta'} \cap h^{-1}(s' - \epsilon)$ and $H_{\theta''} \cap h^{-1}(s'' + \epsilon)$ are one-sheeted hyperboloids.

By the intermediate value theorem, there must be some H_{θ} so that $H_{\theta} \cap h^{-1}(s)$ includes p , and for small epsilon $H_{\theta} \cap h^{-1}(t_0 - \epsilon)$ and $H_{\theta} \cap h^{-1}(t_0 + \epsilon)$ are both one-sheeted hyperboloids near p . Then H_{θ} is a singular 3-manifold. We illustrate this situation in Figure 2.3. (We hope that what just took a lot of words to say becomes obvious in Fig. 2.3.) We see now that it is *essential* that the 3-dimensional fibers of Z^4 “resolve” the same way at cone p .

We make this more precise:

Definition 2.5. Let $\mathcal{F}_t$ be a movie of singular fibrations on a 4-manifold Z^4 with Morse function $h : Z^4 \rightarrow I$. Let t_0 be a regular value of h . Let $x_{t_0} \in h^{-1}(t_0)$ be a critical point of $\mathcal{F}_{t_0}$. Let $B \subset Z^4$ be a small neighborhood of x_{t_0} . Take B small so

	index of p	
	1	2
$T_{\mathcal{F},t_0}(p) > 0$	type I	type II
$T_{\mathcal{F},t_0}(p) < 0$	type II	type I

	index of q	
	0	3
$T_{\mathcal{F},t_0}(q) > 0$	type $\emptyset$	type III
$T_{\mathcal{F},t_0}(q) < 0$	type III	type $\emptyset$

	index of b in $\mathcal{F}_t$ restricted to $\partial h^{-1}(t)$	
	0	2
$T_{\mathcal{F},t_0}(b) > 0$	type I_B	type II_B
$T_{\mathcal{F},t_0}(b) < 0$	type II_B	type I_B

Table 2.1: We define the type of a cone or half-cone p , a dot or half-dot q , or a bowl b in a movie of singular fibrations.

that for some small $\epsilon > 0$, there is a unique critical point x_t of $\mathcal{F}_t$ in the interior of B for $t \in (t_0 - \epsilon, t_0 + \epsilon)$.

Let

$$T_{\mathcal{F},t_0}(x) = \left. \frac{d}{ds} \right|_{s=t_0} \mathcal{F}_s(x_s).$$

Let $p \in \mathcal{F}_{t_0}^{-1}(\theta_p)$ be a cone or half-cone singularity. Let $q \in \mathcal{F}_{t_0}^{-1}(\theta_q)$ be a dot or half-dot singularity. Let $b \in \partial \mathcal{F}_{t_0}^{-1}(\theta_b)$ be a bowl point. Assume $T_{\mathcal{F},t_0}(\theta_p)$, $T_{\mathcal{F},t_0}(\theta_q)$, $T_{\mathcal{F},t_0}(\theta_b) \neq 0$.

We define the type of p or q or b with respect to $\mathcal{F}_t$ in Table 2.1.

See Figure 2.4. In words, if as t decreases the leaves of $\mathcal{F}_t$ near a cone p change from hyperboloids of two sheets to hyperboloids of one sheet (the leaves “fuse together”), we say p is type I. If the leaves change from hyperboloids of one sheet to hyperboloids of two sheets (the leaves “split apart”), we say p is type II. Similarly for half-cones.

If as t decreases the leaves of $\mathcal{F}_t$ near a dot q shrink, then we say q is of type III. If the spheres expand, we say q is of type $\emptyset$. Similarly for half-dots.




Figure 2.4: The type of a singularity p of $\mathcal{F}_t$. The arrow indicates the direction in which a singular cross-section of H_θ is resolved as t decreases.

if as t decreases the leaves of $\mathcal{F}_t$ near a bowl point b move away from $\partial h^{-1}(t)$ near b , then we say b is type II_B . If as t decreases, the leaves of $\mathcal{F}_t$ move toward $\partial h^{-1}(t)$ near b , then we say b is type I_B .

For $H_\theta = \cup_t \mathcal{F}_t^{-1}(\theta)$ to be nonsingular, it must be the case that a singularity of $\mathcal{F}_t$ does not change type as t varies (away from critical points of h).

Remark 2.6. The name of a type of an interior cone or dot refers to a relative handle in $H_\theta = \cup_t \mathcal{F}_t^{-1}(\theta)$ (in a handle decomposition relative to $\mathcal{F}_1^{-1}(\theta)$). A cone of type I or II contributes a 1- or 2-handle, respectively. A dot of type $\emptyset$ or III contributes a 0- or 3-handle, respectively.

A half-cone or half-dot contributes a handle with half the usual attaching region. That is, a copy of $B^k \times B^{3-k}$ attached along (one hemisphere of $S^{k-1} \times B^{3-k}$ (where S^0 is its own hemisphere). Therefore, a half-cone or half-dot of type I or type $\emptyset$ contributes a 1- or 0-handle respectively. A half-cone or half-dot of type II or III do not contribute handles to H_θ .

A bowl point of type II_B contributes a 2-handle to H_θ . A bowl point of type I_B does not contribute a handle to H_θ .

Definition 2.7. Let $\mathcal{F}_t$ be a movie of singular fibrations. We will keep track of the types and indices of singularities as t decreases via a diagram we call a *singularity chart* of $\mathcal{F}_t$. The vertical axis is t (with $t = 1$ at the top and $t = 0$ at the bottom). The horizontal axis is $S^1 = \mathcal{F}_t(h^{-1}(t))$. We plot the singular values of $\mathcal{F}_t$, so that as t varies these points trace out arcs with some endpoints where a singularity is born or dies. We label each arc with the type of the corresponding singularity. Singularities in $\partial h^{-1}(t)$ are drawn with dashed lines. (Note this is essentially a Cerf chart.)

Definition 2.8. A singularity chart is said to be *valid* if it satisfies the following conditions.

- All arcs are smooth in their interiors and are never vertical.
- Arcs may intersect transversely in their interiors. At each endpoint, there are two or three arcs which share that endpoint. Near the endpoint, the chart looks like one of the subcharts in Figure 2.5.

If a singularity chart is not valid, we say it is *invalid*. See Figure 2.6 for an example of a valid singularity chart.

Note that the index of a singularity and the slope of its arc in a singularity chart determine its type. If on the arc, θ decreases as t decreases (i.e. the arc has positive slope), then the index and type agree (e.g. an index-2 cone is type II). If θ increases as t decreases (i.e. the arc has negative slope), then the index and type do not agree (e.g. an index-2 cone is type I). Therefore, labelling each arc of a valid singularity chart with only the type of the singularity it corresponds to also determines the indices of each singularity.

Remark 2.9. In a valid singularity chart, singularity arcs which cross in their interior (i.e. correspond to singular points whose heights interchange) may have the same or different types. Crossing data does not affect the validity of the chart.




Figure 2.5: Possible configurations of two or three arcs meeting at an endpoint in a valid singularity chart.

Definition 2.10. We say that a movie of singular fibrations $\mathcal{F}_t$ is *valid* if the singularity chart for $\mathcal{F}_t$ is valid after potentially perturbing $\mathcal{F}_t$ away from critical points of h .

That is, $\mathcal{F}_t$ is valid if it is valid near critical points of h , and each arc in the singularity chart for $\mathcal{F}_t$ attains one type (sign of slope), but may sometimes have undefined type (vertical slope). Singularity arcs may be tangent or coincide in some interval, but a small perturbation of $\mathcal{F}_t$ should make them intersect transversely.

In the singularity chart for a valid movie, we label each arc with the unique type it ever attains.




Figure 2.6: An example of a valid singularity chart. Note that no arc is ever vertical. Each endpoint locally looks like one of the images in Fig. 2.5. Colors/shades of the arcs are only meant to help the reader view the image; they contain no mathematical meaning.

From now on, when $\mathcal{F}_t$ is a valid movie, we will implicitly perturb $\mathcal{F}_t$ to have valid singularity chart.

Theorem 1.11 essentially follows from the definition of a valid singularity chart.

Theorem 1.11. Let $\mathcal{F}_t$ be a movie of singular fibrations on Z^4 with a valid singularity chart so that $\mathcal{F}_0$ and $\mathcal{F}_1$ are nonsingular. Then Z^4 is fibered over S^1 by $\{H_\theta\}$, where $H_\theta = \cup_t \mathcal{F}_t^{-1}(\theta)$.

Proof. Because the types of an arc in a singularity chart $\mathcal{C}$ for $\mathcal{F}_t$ never change, each H_θ is nonsingular away from points in Z^4 corresponding to endpoints of arcs in $\mathcal{C}$.

Where $\mathcal{C}$ has a cusp (as in a standard Cerf diagram), the H_θ are smooth. Near the endpoint, the arcs are contributing geometrically cancelling handles to each H_θ (Or, in the case of a type II,III- or I_B -,II- boundary pair, no handles at all.)

Where three arcs in $\mathcal{C}$ have a common endpoint (i.e. at a cusp of an interior and boundary singularity, with another boundary singularity on the other side), the H_θ are similarly smooth. Near the endpoint, the interior singularity arc contributes a k -handle, where $k = \{0, 1, 2, 3\}$ if the singularity is type $\{\emptyset, \text{I}, \text{II}, \text{III}\}$. The boundary singularity cancelled by the interior contributes a $D^j \times D^{3-j}$ attached along (hemisphere of $S^{j-1} \times D^{3-j}$, where $j = \{0, 1, 2, 3\}$ if the singularity is type $\{\emptyset, \text{I or } \text{I}_B, \text{II or } \text{II}_B, \text{III}\}$.

Suppose $k = 1$ and $j = 2$. The handle $D^1 \times D^2$ and half-handle $D^2 \times D^1$ glue along (hemisphere of $S^1 \times D^1$ to form a D^3 attached along (hemisphere of $S^0 \times D^2$, exactly the handle contribution of a type I half-cone. (Recall that S^0 is its own hemisphere.) Similarly for other cancelling k and j . These points in the singularity chart should be thought of as a standard critical point birth/death, intersected with a half-ball whose boundary meets the critical point.

Finally, we consider a common endpoint of two arcs in $\mathcal{C}$ which is not a cusp. This endpoint is at a local maximum or minimum of arcs in $\mathcal{C}$, so can be thought of as a birth or death of two arcs. The two arcs are of the same type (opposite index). This pair of births/deaths corresponds to a critical point of h .

Recall in the definition of a singular fibration on a singular 3-manifold $M^3 = h^{-1}(t)$, we specify six possible configurations of $\mathcal{F}^{-1}(t)$ near a singular point in ∂M^3 or $\mathring{M}^3$. (this was Fig. 2.2). Say $x \in h^{-1}(t_0)$ is a critical point of h . Then $\mathcal{F}^{-1}(t_0)$ near x determines $\mathcal{F}^{-1}(t_0 \pm \epsilon)$ near x , up to an automorphism of S^1 .

We assume that $\mathcal{F}_t$ is valid, which further constrains this automorphism of S^1 . This determines the topology of H_θ near x . In particular, the H_θ containing x is nonsingular near x . See Figure 2.7.

Thus, $\{H_\theta\}$ is a (transversely oriented) codimension-1 foliation of Z^4 with compact leaves. That is, $\{H_\theta\}$ is a fibration of Z^4 over S^1 .

□




Figure 2.7: If $\mathcal{F}_t$ has a valid movie of singular fibrations, then near a critical point of h , $\mathcal{F}_t$ looks like one of these movies or their inverses ($t \mapsto 1 - t$). We shade the H_θ that includes the critical point of h , and find that H_θ is smooth near the critical point. We have drawn each picture as if the singularity is in ∂M^3 ; double the picture to obtain a movie containing an interior critical point of h .

Chapter 3

Building blocks: simple movies of fibrations

In this chapter, we construct several explicit movies of singular fibrations.

Every movie $\mathcal{F}_t : Z^4 \rightarrow S^1$ (with height function $h_F : Z^4 \rightarrow I$) from this chapter has the following property: Up to isotopy $\mathcal{F}_t$ agrees with $\mathcal{F}_1$ on $(\partial h^{-1}(1)) \cap (\pi \partial h^{-1}(t))$, under a natural projection π .

Let $\mathcal{G}$ be an arbitrary singular fibration on a 3-manifold M^3 . Suppose that $\mathcal{G}$ agrees with $\mathcal{F}_1$ on some copy $h_F^{-1}(1) \subset M^3$. Then we can extend $\mathcal{G}$ to a movie $\mathcal{G}_t$ on $[(M^3 \setminus h_F^{-1}(1)) \times I] \cup Z^4$ where $\mathcal{G}_t = \mathcal{G}$ outside Z^4 and $\mathcal{G}_t = \mathcal{F}_t$ inside Z^4 .

We will say that we build $\mathcal{G}_t$ by “playing” Movie $\mathcal{F}_t$, where the identification $h_F^{-1}(1) \hookrightarrow M^3$ is clear. We might refer to $\mathcal{G}_t$ as agreeing with the “top” of $\mathcal{F}_t$ in $h_F^{-1}(1) \subset M^3$.

At the end of each section, we produce a valid singularity chart for each movie.

To make the definition of these movies simpler, we will consider a special singular fibration on B^3 .

Definition 3.1. A singular fibration $\mathcal{F}$ on $B^3 = D^2 \times I / (D^2 \times 0 \sim \text{pt}, D^2 \times 1 \sim \text{pt})$ is *standard* if each component of $\mathcal{F}^{-1}(\theta)$ (for all θ) is $D^2 \times \text{pt}$. A standard singular fibration of B^3 has no interior singularities, and has two half-dot singularities in ∂B^3 .

3.1 Adding or cancelling singular points

We define movies corresponding to adding or cancelling a pair of singularities.

Movie 1 (Interior stabilization and destabilization). See Figure 3.1.

Let $Z^4 = B^3 \times I$, where $h : Z^4 \rightarrow I$ is projection onto the second factor. Let $\mathcal{F}_1$ be a standard singular fibration of $B^3 \times 1$. View $\mathcal{F}_1 : B^3 \rightarrow S^1$ as a circular Morse function. As t decreases to $1/2$, obtain $\mathcal{F}_t$ from $\mathcal{F}_1$ by perturbing to add a cancelling-index pair. Then $\mathcal{F}_{1/2}$ agrees with $\mathcal{F}_1$ in a neighborhood of ∂B^3 , but $\mathcal{F}_{1/2}$ has exactly two interior singularities: one cone and one dot if the cancelling pair are index-0,-1 or -2,-3; two cones if the cancelling pair are index-1,-2. Extend $\mathcal{F}_t$ vertically to $t = 0$. We call $\mathcal{F}_t$ an interior stabilization movie (we generally include the indices of the birthed critical points but may drop the word “interior”).

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1 - t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. With respect to $\bar{h}$, we call $\mathcal{G}_t$ an interior destabilization movie.

We note that in the description of $\mathcal{F}_t$, it is not clear what are the types of the two singularities. In fact, we may parametrize $\mathcal{F}_t$ so that the types of the singularities either both agree or both disagree with their indices (i.e. an index-0,-1 pair may be types $\emptyset$ and I or types III and II). See Figure 3.2.

Movie 2 (Boundary stabilization and destabilization). See Figure 3.3.

Let $Z^4 = B^3 \times I$, where $h : Z^4 \rightarrow I$ is projection onto the second factor. Let $\mathcal{F}_1$ be a standard singular fibration of $B^3 \times 1$. View $\mathcal{F}_1 : B^3 \rightarrow S^1$ as a circular Morse function. Fix $x \in \partial h^{-1}(1)$ so that there are no interior or boundary singularities of $\mathcal{F}_1$ in a small neighborhood $\nu(x)$.




Figure 3.1: A diagram of an interior stabilization movie. As t decreases, we perturb $\mathcal{F}_t$ to add a cancelling pair of critical points. Top: an index-0,-1 or -2,-3 pair, depending on the orientation of $\mathcal{F}_t$. Bottom: an index-1,-2 pair (the order depends on the orientation of $\mathcal{F}_t$).




Figure 3.2: Each row is a schematic of an interior stabilization movie (either index-0,-1 or -2,-3). In the top movie, the cone is type I and the dot is type $\emptyset$. In the bottom, the cone is type II and the dot is type III.




Figure 3.3: As t decreases, we stabilize or destabilize $\mathcal{F}_t$ at a point in $\partial h^{-1}(t)$.

View $\mathcal{F}_1 : B^3 \rightarrow S^1$ as a circular Morse function. As t decreases to $1/2$, obtain $\mathcal{F}_t$ from $\mathcal{F}_1$ by perturbing to add a cancelling pair of critical points at x . Then $\mathcal{F}_{1/2}$ agrees with $\mathcal{F}_1$ outside $\nu(x)$, but has two boundary singularities in $\nu(x)$.

If the stabilization is 0-,1- or 2-,3- stabilization, then the new boundary singularities are a half-cone and half-dot. If the stabilization is a 1-,2- stabilization, then the new boundary singularities are a half-cone and a bowl. We call $\mathcal{F}_t$ the boundary stabilization movie (and specify the indices of the birthed singularities).

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1 - t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. With respect to $\bar{h}$, we call $\mathcal{G}_t$ the boundary destabilization movie.

As in the interior stabilization movie, the types of the singularities of $\mathcal{F}_t$ may be taken to agree with their indices, or to be opposite.

In Figure 3.4, we give a valid singularity chart for each interior stabilization and destabilization movie. We give two charts for each, which describe different parametrizations of a (de)stabilization movie. We omit the boundary singularities, which persist through the whole movie.

In Figure 3.5, we give a valid singularity chart for each boundary stabilization and destabilization movie. We give two charts for each, which describe different parametrizations of a (de)stabilization movie. We omit other singularities which persist through the whole movie.




Figure 3.4: Valid singularity charts for each interior stabilization/destabilization. The left half are interior stabilization, while the right half are destabilization. Top row: 0-, 1-(de)stabilization. Middle row: 1-, 2-(de)stabilization. Note that either of the cones may be taken to be the type I cone. Bottom row: 2-, 3-(de)stabilization.




Figure 3.5: Valid singularity charts for each boundary stabilization/destabilization. The left half are boundary stabilization, while the right half are boundary destabilization. Top and second row: 1-, 2- (de)stabilization. Third row: 0-, 1-(de)stabilization. Bottom row: 2-, 3-(de)stabilization.




Figure 3.6: The first boundary fusion movie.

3.2 Movies on Z^4 where $h : Z^4 \rightarrow I$ has one critical point

In the proof of Theorem 1.11, we implicitly described movies of singular fibrations on Z^4 where $h : Z^4 \rightarrow I$ has one critical point. Now we make this more explicit. We start with the case that the critical point of h is contained in ∂Z^4 .

Movie 3 (Boundary fusion/compression movie 1). Let $Z^4 \cong B^4$ and $h : Z^4 \rightarrow I$ so that $h^{-1}(1) = B^3$, $h^{-1}(0) \cong S^1 \times D^2$, and h has one critical point. Say the critical value is $1/2$.

Let $\mathcal{F}_t$ be the movie of Figure 3.6. That is, let $\mathcal{F}_t$ be a standard fibration on B^3 for $t > 1/2$. One leaf of $\mathcal{F}_{1/2}$ is a wedge of two disks along a boundary point. For $t < 1/2$, $\mathcal{F}_t$ has two half-cone singularities of opposite index. $\mathcal{F}_t$ has a singularity chart containing a birth of two type II half-cones, of opposite index.

We call $\mathcal{F}_t$ the first boundary fusion movie (or boundary fusion movie 1).

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1 - t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. With respect to $\bar{h}$, $\mathcal{G}_t$ has a singularity chart containing a death of two type I half-cones, of opposite index.

We call $\mathcal{G}_t$ the first boundary compression movie (or boundary compression movie 1).




Figure 3.7: The second boundary fusion movie. We only draw a neighborhood of the arc between the two bowls, ignoring the other boundary singularities.

Movie 4 (Boundary fusion/compression movie 2). Let $Z^4 \cong B^4$ and $h : Z^4 \rightarrow I$ so that $h^{-1}(1) = B^3$, $h^{-1}(0) \cong S^1 \times D^2$, and h has one critical point. Say the critical value is $1/2$.

Let $\mathcal{F}_t$ be the movie of Figure 3.7. $\mathcal{F}_1$ is a fibration with no interior singularities, two bowls (of opposite indices), two half-cones (opposite indices), and two half-dots (opposite indices). In particular, there is an arc between the two bowl points which is transverse to the leaves of $\mathcal{F}_1$, as pictured. In $\mathcal{F}_{1/2}$, one leaf $\mathcal{F}_{1/2}^{-1}(\theta_0)$ is a disk and contains the critical point of h . For $t < 1/2$, for θ near θ_0 , $\mathcal{F}_t^{-1}(\theta)$ is an annulus.

$\mathcal{F}_t$ has a singularity chart containing a death of two type I_B bowls, of opposite index.

We call $\mathcal{F}_t$ the second boundary fusion movie (or boundary fusion movie 2).

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1 - t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. With respect to $\bar{h}$, $\mathcal{G}_t$ has a singularity chart containing a birth of two type II_B bowls, of opposite index.

We call $\mathcal{G}_t$ the second boundary compression movie (or boundary compression movie 2).

Movie 5 (Boundary fusion/compression movie 3). Let $Z^4 \cong B^4$ and $h : Z^4 \rightarrow I$ so that $h^{-1}(1) = B^3 \sqcup B^3$, $h^{-1}(0) = B^3$, and h has one critical point. Say the critical value is $1/2$.

Let $\mathcal{F}_t$ be the movie of Figure 3.8.




Figure 3.8: The third boundary fusion movie.

That is, let $\mathcal{F}_t$ be a standard fibration on each component of $B^3 \sqcup B^3$ for $t > 1/2$. The leaf of $\mathcal{F}_{1/2}$ containing the critical point of h is a wedge of two disks. For $t < 1/2$, $\mathcal{F}_t$ is a standard fibration.

$\mathcal{F}_t$ has a singularity chart containing a birth of two type I half-cones, of opposite index.

We call $\mathcal{F}_t$ the third boundary fusion movie (or boundary fusion movie 3).

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1 - t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. With respect to $\bar{h}$, $\mathcal{G}_t$ has a singularity chart containing a death of two type II half-cones, of opposite index.

We call $\mathcal{G}_t$ the third boundary compression movie (or boundary compression movie 3).

Movie 6 (Boundary fusion/compression movie 4). Let $Z^4 \cong B^4$ and $h : Z^4 \rightarrow I$ so that $h^{-1}(1) = B^3 \sqcup B^3$, $h^{-1}(0) = B^3$, and h has one critical point. Say the critical value is $1/2$.

Let $\mathcal{F}_t$ be the movie of Figure 3.9.

That is, let $\mathcal{F}_t$ be a standard fibration on each component of $B^3 \sqcup B^3$ for $t > 1/2$. One leaf of $\mathcal{F}_{1/2}$ is the critical point of h . For $t < 1/2$, $\mathcal{F}_t$ is a standard fibration.




Figure 3.9: The fourth boundary fusion movie.

$\mathcal{F}_t$ has a singularity chart containing a death of two type $\emptyset$ half-dots, of opposite index.

We call $\mathcal{F}_t$ the fourth boundary fusion movie (or boundary fusion movie 4).

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1 - t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. With respect to $\bar{h}$, $\mathcal{G}_t$ has a singularity chart containing a birth of two type III half-dots, of opposite index.

We call $\mathcal{G}_t$ the fourth boundary compression movie (or boundary compression movie 4).

Movie 7 (Boundary birth/death movie 1). Let $Z^4 \cong B^4$ and $h : Z^4 \rightarrow I$ so that $h^{-1}(1) = B^3$, $h^{-1}(0) = B^3 \setminus \nu(\text{point}) \cong S^2 \times I$, and h has one critical point. Say the critical value is $1/2$.

Let $\mathcal{F}_t$ be the movie of Figure 3.10.

That is, let $\mathcal{F}_t$ be a standard fibration on each component of $B^3 \sqcup B^3$ for $t > 1/2$. For $t < 1/2$, $\mathcal{F}_t$ agrees with a standard fibration in a neighborhood of ∂B^3 . $\mathcal{F}_t$ has two bowl singularities on the other boundary component of $h^{-1}(t)$.

$\mathcal{F}_t$ has a singularity chart containing a birth of two type I_B bowls, of opposite index.

We call $\mathcal{F}_t$ the first boundary birth movie (or boundary birth movie 1).




Figure 3.10: The first boundary birth movie.




Figure 3.11: The second boundary birth movie.

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1 - t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. With respect to $\bar{h}$, $\mathcal{G}_t$ has a singularity chart containing a death of two type II_B bowls, of opposite index.

We call $\mathcal{G}_t$ the first boundary death movie (or boundary death movie 1).

Movie 8 (Boundary birth/death movie 2). Let $Z^4 \cong B^4$ and $h : Z^4 \rightarrow I$ so that $h^{-1}(1) = \emptyset$, $h^{-1}(0) = B^3$, and h has one critical point. Say the critical value is $1/2$.

Let $\mathcal{F}_t$ be the movie of Figure 3.11.

That is, let $\mathcal{F}_t$ be a standard fibration on B^3 for $t < 1/2$. $\mathcal{F}_{1/2}$ maps $h^{-1}(1/2) = \text{pt}$ to some value of θ .

$\mathcal{F}_t$ has a singularity chart containing a birth of two type $\emptyset$ half-dots, of opposite index.

We call $\mathcal{F}_t$ the second boundary birth movie (or boundary birth movie 2).

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1 - t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. With respect to $\bar{h}$, $\mathcal{G}_t$ has a singularity chart containing a death of two type III half-dots, of opposite index.

We call $\mathcal{G}_t$ the second boundary death movie (or boundary death movie 2).

Movie 9. Let $h : Z^4 \rightarrow I$ be Morse and $\mathcal{F}_t$ on Z^4 be a boundary fusion, compression, birth, or death movie. Let $x \in \partial Z^4$ be the critical point of h . Let $X^4 = Z^4 \cup_{\partial Z^4 \cap \nu(x)} \overline{Z^4}$ be a double of Z^4 along x , and $\tilde{h} : X^4 \rightarrow I$ be given by $\tilde{h} = h \cup \bar{h}$.

Let $\mathcal{G}_t$ be a movie of singular fibrations on X^4 given by $\mathcal{G}_t = \mathcal{F}_t \cup \overline{\mathcal{F}_t}$ (that is, $\mathcal{G}_t$ agrees with $\mathcal{F}_t$ on Z^4 , and agrees with $\overline{\mathcal{F}_t}$ on $\overline{Z^4}$).

We call $\mathcal{G}_t$ an *interior singularity movie*. Let $\mathcal{C}_{\mathcal{F}}$ and $\mathcal{C}_{\mathcal{G}}$ be valid charts for $\mathcal{F}$ and $\mathcal{G}$, respectively. We give a table with the name of $\mathcal{G}_t$ and a description of part of $\mathcal{C}_{\mathcal{F}}$ and $\mathcal{C}_{\mathcal{G}}$.

$\mathcal{F}_t$	$\mathcal{G}_t$	in $\mathcal{C}_{\mathcal{F}}$ (boundary)	in $\mathcal{C}_{\mathcal{G}}$ (as interior)	index of c.p. of h
Boundary fusion 1	Index-2 birth	II birth	II birth	2
Boundary fusion 2	Index-1 death	I_B death	I death	2
Boundary fusion 3	First index-1 birth	I birth	I birth	3
Boundary fusion 4	Index-0 death	$\emptyset$ death	$\emptyset$ death	3
Boundary birth 1	Second Index-2 birth	I_B birth	I birth	2
Boundary birth 2	Index-0 birth	$\emptyset$ birth	$\emptyset$ birth	4
Boundary compression 1	Index-1 death	I death	I death	2
Boundary compression 2	Index-2 birth	II_B birth	II birth	2
Boundary compression 3	First Index-2 death	II death	II death	1
Boundary compression 4	Index-3 birth	III birth	III birth	1
Boundary death 1	Second Index-1 death	II_B death	II death	2
Boundary death 2	Index-3 death	III death	III death	0

The names of these movies may seem unwieldy, but the logic is for the name of $\mathcal{G}_t$ to simply describe the critical points introduced in $\mathcal{G}_t$ as t decreases from 1 to 0. An index- k birth corresponds to an index- $(4 - k)$ critical point of the ambient height function, while an index- k death corresponds to an index- $(3 - k)$ critical point of the ambient height function. (Again, there is some confusion as we generally work from top to bottom, but this viewpoint is convenient for this paper.)

In Figure 3.12, we give valid singularity charts for the boundary fusion, compression, birth, death movies and the interior singularity movies, omitting the persisting singularities.




Figure 3.12: Valid singularity charts for the boundary fusion, compression, birth, death and interior singularity movies.

3.3 Positioning singularities in a fibration

In this section, we define movies $\mathcal{F}_t$ on $M^3 \times I$ which allow us to “position” the singularities of $\mathcal{F}_1$, in the sense of choosing the singular values of $\mathcal{F}_0$ (with some constraints). Imprecisely, both of these movies essentially “flow” some choice of leaves of $\mathcal{F}_1$ through a singularity. These movies exchange the heights of critical points of $\mathcal{F}_t$ as t decreases.

Movie 10 (Positioning a dot or half-dot along an arc). Let $\mathcal{F}_1$ be a singular fibration on M^3 . Suppose q is a cone point, and p is a dot or half-dot of $\mathcal{F}_1$. Assume there exists an injective $\gamma : [0, 1] \rightarrow M^3$ transverse to the leaves of $\mathcal{F}_1$ so that $\gamma(0) = q$, $\gamma(1) = p$.

If $p \in \mathring{M}^3$, then assume $\gamma(I) \subset \mathring{M}^3$ and the leaf component of $\mathcal{F}_1$ containing $\gamma(t)$ is a 2-sphere for $0 < t < 1$.

If $p \in \partial M^3$, then assume $\gamma(I) \cap \partial M^3 = p$ and the leaf component of $\mathcal{F}_1$ containing $\gamma(t)$ is a 2-hemisphere for $0 < t < 1$.

Fix $\epsilon > 0$ small, and let $\eta : [0, 1] \rightarrow M^3$ be given by $\eta(s) = \gamma((1 - \epsilon)s + \epsilon)$.

First assume p is in the interior of $h^{-1}(t)$ (so p is a dot). We define a movie $\mathcal{F}_t$ on $M^3 \times I$ as follows:

- If $x \in M^3$ is not in a leaf component of $\mathcal{F}_1$ meeting $\eta([0, 1])$, then $\mathcal{F}_t(x) = \mathcal{F}_1(x)$ for all t .
- Let B be the set on which we have not defined $\mathcal{F}_t$. If p is a dot singularity (in the interior of $h^{-1}(t)$), write $\overline{B} = S^2 \times I / (S^2 \times 1 \sim \text{pt})$ so $p = S^2 \times 1 / \sim$, and $\mathcal{F}_1(S^2 \times s) = \mathcal{F}_1(\eta(s))$.
- If p is a half-dot, write $B = D^2 \times I / (D^2 \times 1 \sim \text{pt})$ so $p = D^2 \times 1 / \sim$, and $\mathcal{F}_1(D^2 \times s) = \mathcal{F}_1(\gamma(s))$.
- Define $\mathcal{F}_t$ by

$$\mathcal{F}_t(S^2 \times s) = \mathcal{F}_1(\eta(ts + (1-t)\epsilon s)).$$

As t decreases, we are shrinking the range of $\mathcal{F}_t|_B$ while fixing $\mathcal{F}_t|_{M^3 \setminus B}$. See Figure 3.13 for an illustration.

Note that when positioning a dot or half-dot, the (components of) the level sets of $\mathcal{F}_t$ are identical for all t . In the beginning of this section, we said this movie would “position” a singularity. The singularities $p, q \in M^3$ of $\mathcal{F}_1$ are also singularities of $\mathcal{F}_0$; we have not moved the singular points within M^3 . However, in $M^3 \times 0$ the arc γ is very short in the sense that $\mathcal{F}_0(\gamma(I))$ is very short in S^1 .

If $\epsilon > 0$ is taken to be sufficiently small, then $\gamma((0, 1))$ does not intersect any singular leaf of $\mathcal{F}_0$.

Movie 11 (Positioning a cone along an arc). Consult Figure 3.14. This image conveys the idea of this movie better than the ensuing text.

Let $\mathcal{F}_1$ be a singular fibration on M^3 . Suppose q is a cone point of $\mathcal{F}_1$. Let $\gamma : [-1, 1] \rightarrow \mathring{M}^3$ be injective so that $\gamma(0) = q$, $\mathcal{F}_1(\gamma(s)) = \mathcal{F}_1(\gamma(-s))$ for all s , and




Figure 3.13: Diagram of positioning a dot or half-dot. We draw a neighborhood of B , which might be a solid torus or a ball. On the light gray leaf components (and all components not illustrated), $\mathcal{F}_t$ is fixed for all t . As t decreases, the range of $\mathcal{F}_t(B)$ contracts.

$\gamma(I)$ is transverse to the leaves of $\mathcal{F}_1$ and does not meet any singularities except q . Let B be a small ball containing $\gamma(I)$. Choose γ so that near q , $\gamma(\pm\epsilon)$ meets the two-sheeted hyperboloid leaves of $\mathcal{F}_1|_B$ (rather than the one-sheeted leaves).

We define the movie $\mathcal{F}_t$ by:

- If x is not in $\mathring{B}$, then $\mathcal{F}_t(x) = \mathcal{F}_1(x)$ for all t .
- The leaves of $\mathcal{F}_t|_B$ are (up to isotopy) hyperboloids, except for one cone.
- At time t , the cone of $\mathcal{F}_t|_B$ lies in $\mathcal{F}_t^{-1}\mathcal{F}_1(\gamma(1-t))$. The leaves of $\mathcal{F}_1|_B$ meeting $\gamma(s)$ for $|s| > 1-t$ are still two-sheeted hyperboloid leaves of $\mathcal{F}_t|_B$, while the leaves of $\mathcal{F}_1$ meeting $\gamma(s)$ for $|s| < 1-t$ become one-sheeted hyperboloid leaves of $\mathcal{F}_t|_B$.

The purpose of this movie is that in $\mathcal{F}_0$, the cone singularity q now lies in $\mathcal{F}_0^{-1}(\mathcal{F}_1(\gamma(\pm 1)))$.




Figure 3.14: Diagram of positioning a cone along an arc γ . We draw a neighborhood of γ . As t decreases, the value of $\mathcal{F}_t(\text{cone point})$ changes from $\mathcal{F}_1(\gamma(0))$ to $\mathcal{F}_1(\gamma(\pm 1))$.

As described, a chart for $\mathcal{F}_t$ has a cone of type I. We will now reparametrize $\mathcal{F}_t$ so that the cone is type II. Let B be a small neighborhood of $\gamma([-1, 1])$ so that q is the only singularity of $\mathcal{F}_1$ contained in B . Let $f : S^1 \rightarrow [0, m]$ be a narrow bump function on $\mathcal{F}_1(q)$. If q is index-1, let $f := -f$.

Let $g_t : S^1 \rightarrow S^1$ be a smooth family of automorphisms with $g_1 = \text{id}$ and $dg_t/dt = -f$. Obtain a movie of singular fibrations $\tilde{\mathcal{G}}_t$ on $B \times I$ defined by $\tilde{G}_t(x) = g_t \circ \mathcal{F}_t(x)$ for $x \in h^{-1}(t)$. If m is taken to be sufficiently large, then the repositioned cone is type II in $\tilde{\mathcal{G}}_t$. Now isotope $\tilde{\mathcal{G}}_t$ near ∂B so that $\tilde{\mathcal{G}}_t|_{\nu(\partial B)}$ agrees with $\mathcal{F}_t$. Let $\mathcal{G}_t$ be a movie on $M^3 \times I$ given by $\mathcal{G}_t(x) = \begin{cases} \tilde{\mathcal{G}}_t(x) & | x \in B \\ \mathcal{F}_t(x) & | x \notin B \end{cases}$. The movies $\mathcal{G}_t$ and $\mathcal{F}_t$ agree near every singularity of $\mathcal{F}_t$ except q , so their charts are identical except for the arc of q , which has opposite slope (and hence opposite type) in the other chart. See Figure 3.15.

Movie 12 (Positioning a cone along a disk). This movie is the inverse of positioning a cone along an arc. However, we wish to write this movie out explicitly.




Figure 3.15: When positioning a cone along an arc γ , suppose γ meets no other singular leaf. By reparametrizing $\mathcal{F}_t$, we find a movie $\mathcal{G}_t$ in which the positioned cone is type II. In bold: $\mathcal{G}_t^{-1}(\theta_0)$ near the positioned cone, as t decreases.

Let $\mathcal{F}_1$ be a singular fibration on M^3 . Suppose q is a cone point of $\mathcal{F}_1$. Let D be an improperly embedded disk in $\mathring{M}^3$ intersecting q , with ∂D contained in one leaf of $\mathcal{F}_1$. Parameterize D so $D = S^1 \times I / (S^1 \times 1 \sim \text{pt})$, where $S^1 \times 0 = \partial D$ and $S^1 \times 1 / \sim = q$. Assume that each $S^1 \times r$ lives in a single leaf of $\mathcal{F}_1$, which intersects D transversely for $0 < r < 1$. Let B be an open neighborhood of D .

We define the movie $\mathcal{F}_t$ by:

- If x is not in B , then $\mathcal{F}_t(x) = \mathcal{F}_1(x)$ for all t .
- The leaves of $\mathcal{F}_t|_B$ are (up to isotopy) hyperboloids, except one cone.
- At time t , the cone of $\mathcal{F}_t|_B$ lies in $\mathcal{F}_t^{-1}\mathcal{F}_1(S^1 \times t)$. The leaves of $\mathcal{F}_1|_B$ meeting D in $S^1 \times r$ are still one-sheeted hyperboloid leafs of $\mathcal{F}_t|_B$ when $r < t$. When $r > 1$, they become two-sheeted hyperboloid leafs of $\mathcal{F}_t|_B$.

The purpose of this movie is that in $\mathcal{F}_0$, the cone singularity q now lies in $\mathcal{F}_0^{-1}(\mathcal{F}_1(\partial D))$.




Figure 3.16: Valid singularity charts for the positioning movies. Each chart contains one arc (we omit the arcs of the singularities we are not positioning).

As described, the repositioned cone is type II in $\mathcal{F}_t$.

Similarly to the “positioning a cone along an arc” movie, we may reparametrize this movie to find a chart in which the repositioned cone is type I (by first reparametrizing in a small neighborhood of $D \times I$ and then extending to $M^3 \times I$).

In Figure 3.16, we give singularity charts for the singularity positioning movies.

3.4 Cancelling interior and boundary singularities

Movie 13 (Cancelling a cone with a half-cone). See Figure 3.17 (top).

Let $\mathcal{F}_1$ be a singular fibration on B^3 which has two half-cone singularities p_1, p_2 on ∂B^3 and one cone q in the interior of B^3 . (Necessarily, the half-cones are both of index i while the cone is index $3 - i$.)

Let $D = S^1 \times I / (S^1 \times 1 \sim \text{pt})$ be a disk improperly embedded in B^3 so that $D \cap p_2 = \emptyset$, $D \cap \partial B^3 = p_1 \in S^1 \times 0$, $q = S^1 \times 1$, D is transverse to the leaves of $\mathcal{F}$ away from p_1 and q and $S^1 \times s$ is contained in a single leaf for each s . We proceed as if positioning q along D . That is, obtain $\mathcal{F}_t$ by perturbing $\mathcal{F}_1$ in a neighborhood of D

so that the cone lies in $\mathcal{F}_t^{-1}(\mathcal{F}_t(S^1 \times t))$. As t grows small, take the cone point to be very close to p_1 . At $t = 0$, the cone meets p_1 and becomes a half-cone of index $3 - i$.

As written, the interior cone of $\mathcal{F}_t$ is type II while the cancelled boundary half-cone is type I. Now we describe a movie $\mathcal{G}_t$ with the same level sets, in which the types are reversed.

Let $\mathcal{G}_1 = \mathcal{F}_1$. Let $D' = [0, \pi] \times I / ([0, \pi] \times 1 \sim \text{pt})$ be a disk improperly embedded in B^3 so that $D' \cap p_2 = \emptyset$, $p_1 = [0, \pi] \times 1$, $q \in [0, \pi] \times 0$, $D' \cap \partial B^3 = \{0, \pi\} \times I$, D' is transverse to the leaves of $\mathcal{F}$ away from p_1 and q and $[0, \pi] \times s$ is contained in a single leaf for each s . We proceed as if positioning p_1 along D' . That is, obtain $\mathcal{G}_t$ by perturbing $\mathcal{G}_1$ in a neighborhood of D' so that the half-cone lies in $\mathcal{G}_t^{-1}(\mathcal{F}_t([0, \pi] \times t))$. As t grows small, take the cone point to be very close to p_1 . At $t = 0$, the cone meets the boundary and becomes a half-cone of index $3 - i$.

We say that $\mathcal{F}_t$ and $\mathcal{G}_t$ are movies of a cone, half-cone death. Turning $\mathcal{F}_t$ or $\mathcal{G}_t$ upside-down yields a movie of a cone, half-cone birth.

Movie 14 (Cancelling a cone with a half-dot). See Figure 3.17 (bottom).

Let $\mathcal{F}_1$ be a singular fibration on M^3 . Suppose q is a cone point of $\mathcal{F}_1$. Let $\gamma : [0, 1] \rightarrow M^3$ so that $\gamma(0) = q$, $\gamma(1)$ is a half-dot p , and $\gamma(I) \cap \partial M^3 = \emptyset$. Assume $\gamma((0, 1))$ is transverse to the leaves of $\mathcal{F}_1$ and does not meet any singular leaf components of $\mathcal{F}_1$. Choose γ so that near q , $\gamma(\epsilon)$ meets the two-sheeted hyperboloid leaves of $\mathcal{F}_1|_{\nu(q)}$ (rather than the one-sheeted leaves).

From $t = 1$ to $1/2$, position the half-dot along γ so that $\gamma((0, 1))$ does not meet any singular leaves of $\mathcal{F}_{1/2}$. Extend γ to an interval $[-\delta, 1]$ so that $\mathcal{F}_{1/2}(\gamma(s)) = \mathcal{F}_{1/2}(\gamma(-s))$, as in the “positioning a cone along an arc” movie.

Let B be a small neighborhood of $\gamma((-\delta, 1))$.

We define the movie $\mathcal{F}_t$ by:

- If x is not in $\mathring{B}$, then $\mathcal{F}_t(x) = \mathcal{F}_1(x)$ for all t .




Figure 3.17: Top: a schematic of the cone, half-cone death. Bottom: a cone, half-dot death.

- The leaves of $\mathcal{F}_t|_B$ are (up to isotopy) hyperboloids, except one cone.
- At time t , the cone of $\mathcal{F}_t|_B$ lies in $\mathcal{F}_t^{-1}\mathcal{F}_1(\gamma(1-t))$. The leaves of $\mathcal{F}_1|_B$ meeting $\gamma(s)$ for $|s| < t$ are still two-sheeted hyperboloid leaves of $\mathcal{F}_t|_B$, while the leaves of $\mathcal{F}_1$ meeting $\gamma(s)$ for $|s| > t > 0$ become one-sheeted hyperboloid leaves of $\mathcal{F}_t|_B$.

As t grows small, take the cone to be very close to p . At $t = 0$, the cone meets p and becomes a bowl singularity.

As described, the cone is type I and the half-dot is type $\emptyset$. Because $\gamma((-\delta, 1))$ does not meet any singular leaves of $\mathcal{F}_{1/2}$, we can reparametrize $\mathcal{F}_t$ to find a movie $\mathcal{G}_t$ with the same level sets in which the positioned cone is type II (and the half-dot is type III) as follows:

- Let $f : S^1 \rightarrow [0, m]$ be a narrow bump function on $\mathcal{F}_{1/2}(\gamma([-\delta, 1]))$. If q is index-1, let $f := -f$.




Figure 3.18: Top row: valid singularity charts for the cone, half-cone and cone, half-dot deaths. Bottom row: valid singularity charts for the cone, half-cone and cone, half-dot births. We omit the arc of the half-cone or half-dot which is not cancelled or born during the movie.

- Let $g_t : S^1 \rightarrow S^1$ be a smooth family of automorphisms with $g_1 = \text{id}$ and $dg_t/dt = -f$.
- Let $\mathcal{G}_t(x) = g_t \circ \mathcal{F}_t(x)$ for $x \in h^{-1}(t)$.

If m is taken to be sufficiently large, then the repositioned cone is type II in $\mathcal{G}_t$, and the half-dot is type III.

We say that $\mathcal{F}_t$ is a movie of a cone, half-dot death. Turning $\mathcal{F}_t$ upside-down yields a movie of a cone, half-dot birth.

In Figure 3.18, we give a valid singularity chart for the interior and boundary singularity cancellation movies.

3.5 Movies on $B^3 \times I \setminus \nu(\text{saddle, minimum, or maximum disk})$

Now we consider slightly more interesting 4-manifolds. These movies are defined on $Z^4 \cong (B^3 \times I) \setminus (\text{surface with a single singular cross-section})$. Most of the interesting

information of these singular movies is determined by the topology of Z^4 , while (very informally) $\mathcal{F}_t$ does not change much with t .

Movie 15 (Saddle movie). Consult Figure 3.19. Let $h : B^3 \times I \rightarrow I$ be projection onto the second factor. Let $R \subset B^3 \times I$ be a saddle. That is, R is a properly embedded disk with $R \cap (B^3 \times 0) \cong R \cap (B^3 \times 1) \cong \{2 \text{ arcs}\}$ and $h|_R$ is Morse with single index-1 critical point and no index-0 or -2 critical points.

Let $Z^4 = (B^3 \times I) \setminus \nu(R)$ and restrict h to Z^4 . There are two values of t for which $h^{-1}(t)$ is a singular 3-manifold; say these values are $t = 3/4$ and $t = 1/4$. We define a movie of singular fibrations $\mathcal{F}_t : h^{-1}(t) \rightarrow S^1$ on Z^4 as follows:

- Figure 3.19 (top left): Let $\mathcal{F}_1$ be as indicated. There are two half-cone singularities of $\mathcal{F}_1$ in $\partial h^{-1}(1)$, which are both far from $\partial\nu(R)$. The nonsingular leaves of $\mathcal{F}_1$ are either: a single disk, a disjoint union of two disks, or an annulus. There is one cone singularity of $\mathcal{F}_1$ in the interior of $h^{-1}(1)$ corresponding to the compression of the annular leaves. There are no dot singularities.
- Figure 3.19 (top right): At $t = 3/4$, $h^{-1}(3/4)$ is a singular 3-manifold.
- Figure 3.19 (middle left): For $1/4 < t < 3/4$, $h^{-1}(t)$ is a solid of genus-3. $\mathcal{F}_t$ has no singularities in the interior of $h^{-1}(t)$. (The cone leaf at $t > 3/4$ becomes a half-cone meeting $\partial\nu(R)$.) There are four half-cone singularities of $\mathcal{F}_t$ in $\partial h^{-1}(t)$, two of which lie on $\partial\nu(R)$.
- Figure 3.19 (middle right): Again, $h^{-1}(1/4)$ is a singular 3-manifold. Note $h^{-1}(1/4) \cong h^{-1}(3/4)$. Under a rotation, the singular fibration $\mathcal{F}_{1/4}$ is identical to the singular fibration $\mathcal{F}_{3/4}$.
- Figure 3.19 (bottom left): As t decreases to 0, we take $\mathcal{F}_t = \mathcal{F}_{1-t}$ under a rotation so that the movies $\mathcal{F}_t$ (t decreasing from $1/4$ to 0) and $\mathcal{F}_t$ (t increasing from $3/4$ to 1) are identical (under a rotation). This results in a singular




Figure 3.19: Diagram of a saddle movie. This is a movie of singular fibrations on $(B^3 \times I) \setminus \nu(\text{saddle})$.

fibration $\mathcal{F}_0$ with two half-cones on $\partial h^{-1}(0)$ and one cone in the interior of $h^{-1}(0)$.

Note that the cone singularities in $\mathcal{F}_0$ and $\mathcal{F}_1$ are contained in different $\mathcal{F}_t^{-1}(\theta)$'s (different values of θ).

Movie 16 (Minimum/maximum movie). Consult Figure 3.20. Let $h : B^3 \times I \rightarrow I$ be projection onto the second factor. Let $R \subset B^3 \times I$ be a disk with unknotted boundary in $B^3 \times 1$, so that $h|_R$ is Morse with one critical point (necessarily of index 0). Note $R \cap (B^3 \times 0) = \emptyset$.

Let $Z^4 = (B^3 \times I) \setminus \nu(R)$ and restrict h to Z^4 . There are two values of t for which $h^{-1}(t)$ is a singular 3-manifold; say these values are $t = 2/3$ and $t = 1/6$. We define a movie of singular fibrations $\mathcal{F}_t : h^{-1}(t) \rightarrow S^1$ on Z^4 as follows:

- Figure 3.20 (top left): Let $\mathcal{F}_1$ be as indicated. There are two dot singularities in the foliation $\mathcal{F}_1$ induces on $\partial h^{-1}(1)$, which both lie in ∂B^3 . The nonsingular leaves of $\mathcal{F}_1$ are disks and annuli. $\mathcal{F}_1$ has no singularities in the interior of $h^{-1}(1)$.
- Figure 3.20 (top right): As t decreases to $5/6$, play the fourth boundary compression movie.
- Figure 3.20 (left, second row): There are two half-dot singularities on the now spherical boundary component.
- Figure 3.20 (right, second row): Play an interior 0-,1- stabilization and a 2-,3- stabilization near the half-dots. (Do the 0-,1- stabilization near the index-0 half-dot.)
- Figure 3.20 (left, third row): Play a cone, half-dot death movie near each half-dot. Now there are two bowl singularities.




Figure 3.20: Diagram of a minimum movie. This is a movie of singular fibrations on $(B^3 \times I) \setminus \nu(\text{disk with one minimum})$.

- Figure 3.20 (right, third row): Play the first boundary death movie.
- Figure 3.20 (bottom left): As t decreases to 0, we have $h^{-1}(t) = B^3$. There are two dot singularities in the interior of B^3 .

Let $\bar{h} : Z^4 \rightarrow I$ be given by $\bar{h}(t) = h(1-t)$. Let $\mathcal{G}_t = \mathcal{F}_{1-t}$. Now $Z^4 = B^3 \times I \setminus \nu(\text{disk with one maximum with respect to } \bar{h})$. With respect to $\bar{h}$, we call $\mathcal{G}_t$ a maximum movie.

In Figure 3.21, we give a valid singularity chart for the saddle, minimum, and maximum movies.




Figure 3.21: Top row: two singularity charts for the saddle movies (the difference is the indices of the singular points). Bottom left: a chart for the minimum movie. Bottom right: a chart for the maximum movie. The colors are just meant to help the reader distinguish intersecting arcs. We omit arcs corresponding to singularities which persist from $t = 1$ to $t = 0$.

3.6 Composite movies

In this section, we will combine previous movies to define more complicated movies of singular fibrations.

Movie 17 (Simple cancellation movie). Here, we combine the positioning movies with the interior destabilization movie to cancel a cone and dot singularity, when the cone is the nearest singularity of $\mathcal{F}_1$ to the dot.

Let $Z^4 = M^3 \times I$, and $h : M^3 \times I$ be projection onto the second factor. Let $\mathcal{F}_1$ be a singular fibration of $M^3 \times 1$ including a cone singularity q and a dot singularity p , of cancelling indices.

Suppose there exists an arc $\gamma : [0, 1]$ which is transverse to the nonsingular leaves of $\mathcal{F}_1$ so that $\gamma(0) = q$ and $\gamma(1) = p$. Let L be a nonsingular leaf near q so that $\nu(q) \cap L$ has two components. Assume that these two components are in *distinct*




Figure 3.22: If we attempt to cancel a dot with a cone where the two sheets of $\mathcal{F}_t^{-1}(\theta)$ near the cone are in the same component of $\mathcal{F}_t^{-1}(\theta)$, then we cannot actually destabilize the circular Morse function $\mathcal{F}_t$. Left: a local picture of a dot and cone singularity; the two sheets of a leaf near the cone are in the same component of the leaf. Right: We attempt to cancel the dot and cone singularities and create a Reeb component, which is not allowed in the definition of a singular fibration.

components of L . **Assume moreover that $\gamma((0, 1))$ does not meet any singular leaf components of $\mathcal{F}_1$.**

From $t = 1$ to $t = 1/2$, position the dot p along γ so that $\gamma((0, 1))$ intersects no singular leaves of $\mathcal{F}_{1/2}$. Now from $t = 1/2$ to $t = 0$, cancel the singularities (this is an interior destabilization movie).

In order to cancel the singularities, it must be the case that the two components of $L \cap \nu(P)$ are not in the same component of L . See Figure 3.22 for an illustration.

Movie 18 (Generalized cancellation movie). Here, we combine the positioning movies with the interior destabilization movie to cancel a cone and dot singularity which may initially be far apart in $\mathcal{F}_1$.

Let $Z^4 = M^3 \times I$, and $h : M^3 \times I$ be projection onto the second factor. Let $\mathcal{F}_1$ be a singular fibration of $M^3 \times 1$ including a cone singularity q and a dot singularity p , of cancelling indices.

Suppose there exists an arc $\gamma : [0, 1]$ which is transverse to the nonsingular leaves of $\mathcal{F}_1$ so that $\gamma(0) = q$ and $\gamma(1) = p$. Let L be a nonsingular leaf near q so that

$\nu(q) \cap L$ has two components. Assume that these two components are in *distinct* components of L . **We allow $\gamma((0, 1))$ to meet other singular leaf components of $\mathcal{F}_1$, but take $\gamma((0, 1))$ to be far from the actual singularities.**

Note that if M^3 is a connected manifold with nonempty boundary and p is a dot singularity of $\mathcal{F}_1$, then there *must* be a cone q and arc γ satisfying the above setup.

Extend γ to $\gamma : [-1, 1] \rightarrow M^3$ so that $\mathcal{F}_1(\gamma(s)) = \mathcal{F}_1(\gamma(-s))$ for all $s \in [0, 1]$ and γ is still transverse to all the leaves of M^3 (do this by extending γ perpendicularly to the leaves of $\mathcal{F}_1$. If this terminates before extending all the way to -1 , then the extension terminates at a dot $p' = \gamma(-s_0)$. Exchange the roles of p and p' , and let $\gamma(s) := \gamma(-s/s_0)$.)

From $t = 1$ to $t = 2/3$, position the cone q along $\gamma|_{(-1+\epsilon, 1-\epsilon)}$ so that q lies on $\gamma(1 - \epsilon)$, so p and q are very close in $h^{-1}(2/3)$. From $t = 2/3$ to $1/3$, position the dot p along $\gamma((1 - \epsilon, 1))$ so that $\mathcal{F}_{1/3}(\gamma(1 - \epsilon, 1))$ is a very short arc. From $t = 1/3$ to $t = 0$, cancel the singularities (this is an interior destabilization movie).

Remark 3.2. If M^3 is a connected 3-manifold with nonempty boundary and $\mathcal{F}_1$ is a singular fibration on M^3 with $n \geq 1$ dot singularities and r cone singularities, then the generalized cancellation movie allows us to extend $\mathcal{F}_1$ to a valid movie $\mathcal{F}_t$ on $M^3 \times I$ so that $\mathcal{F}_0$ has $n - 1$ dots and $r - 1$ cones. The types of the singularities in $\mathcal{F}_t$ may be chosen freely, except that the cancelled pair of singularities must have cancelling types ($\emptyset$ -, I- or II-, III-).

In particular, if $\mathcal{F}_1$ is a singular fibration on a connected 3-manifold M^3 with nonempty boundary and $\mathcal{F}_1$ has n dot and n cone singularities that are all of types II or III (or all of types $\emptyset$ or I), then $\mathcal{F}_1$ may be extended to a valid movie $\mathcal{F}_t$ on $M^3 \times I$ so that $\mathcal{F}_0$ is nonsingular.

Now we define our last two movies: the band and disk movies. These two movies are more complicated than previous ones, but viewing them as composites allows us to better understand them.

Movie 19 (Band movie). As in the saddle movie, let $h : B^3 \times I \rightarrow I$ be projection onto the second factor. Let $R \subset B^3 \times I$ be a saddle. That is, R is a properly embedded disk with $R \cap (B^3 \times 0) \cong R \cap (B^3 \times 1) \cong (2 \text{ arcs})$ and $h|_R$ is Morse with single index-1 critical point and no index-0 or -2 critical points. Let $b \cong I \times [-\epsilon, \epsilon]$ be a band projected to $B^3 \times 1$ with $S^0 \times [-\epsilon, \epsilon] \subset \partial b$ in $\partial(B^3 \setminus \nu(R))$ so that the band b describes the saddle of R . (That is, b is obtained by perturbing R so the saddle is degenerate and then projecting a neighborhood of the degeneracy to $B^3 \times 1$.) Let $\eta = I \times \text{pt} \subset b$ be a core arc of b .

Restrict h to $(B^3 \times I) \setminus \nu(R)$. Let $\mathcal{G}_1$ be any singular fibration on $h^{-1}(1)$ with the following properties.

- Each leaf of $\mathcal{G}_1$ intersects $(\partial\nu(R)) \setminus \partial B^3$ in arcs.
- Each leaf of $\mathcal{G}_1$ intersects $b \times I \times [-\epsilon, \epsilon]$ in copies of $\text{pt} \times [-\epsilon, \epsilon]$.
- Each intersection of a leaf of $\mathcal{G}_1$ with the interior of b is transverse.

Now consult Figure 3.23. We describe each frame, the arrows indicating decreasing t :

- Top left: A neighborhood B of $\eta(1)$ in $h^{-1}(1)$. Play an interior 1-,2-stabilization movie in B , during the time span $t = 1$ to $1 - \epsilon$. Let q be the cone point of the cone oriented horizontally (opening parallel to the band b); take q to be type I.
- Top right: Take $\gamma : [-1, 1]$ to be an arc in $h^{-1}(\epsilon)$ with $\gamma(-1)$ close to $\eta(0)$, $\gamma(1)$ close to $\eta(1)$, and $\gamma(0) = q$. The arc γ is parallel to the band away from $\eta(1)$, and then near $\eta(1)$ spirals around the end of the band so that we may take $\gamma(s) = \gamma(-s)$ for all s .
- Middle left: Position the cone q along the arc γ . This is a picture of a singular fibration partway through this positioning movie.




Figure 3.23: Some leaves of $\mathcal{G}_t$ (a band movie) near a neighborhood of band b . Outside this neighborhood, $\mathcal{G}_t = \mathcal{G}_1$ for all t .

- Middle right: Isotope $(B^3 \times I) \setminus \nu(R)$ to shrink the band b . We could do this isotopy strictly after the positioning movie, but we believe it is easier to visualize at this step.
- Bottom left: We have finished positioning the cone. Say $t = \epsilon$. There is a set B' containing $\eta(0)$ in $h^{-1}(\epsilon)$ so that in B' , $\mathcal{G}_\epsilon$ agrees with $\mathcal{F}_1$ of the saddle movie. Play the saddle movie in the time span $t = \epsilon$ to 0.
- Bottom right: We obtain a singular fibration $\mathcal{G}_0$ of $h^{-1}(0)$. Note $\mathcal{G}_0$ has exactly two singularities interior to $h^{-1}(0)$ (both are type II cones).




Figure 3.24: Left to right, top to bottom: One leaf in a band movie $\mathcal{G}_t$ as t decreases from 1 to 0.




Figure 3.25: Left to right, top to bottom: Three leaves in a band movie $\mathcal{G}_t$ as t decreases from 1 to 0.

This movie is comparatively complicated, so we include some extra cartoons of leaves in $\mathcal{G}_t$ for decreasing t . See Figures 3.24 and 3.25. In Figure 3.26, we give a more detailed image of the cone singularities in $\mathcal{G}_0$.

Movie 20 (Disk movie). As in the minimum movie, let $h : B^3 \times I \rightarrow I$ be projection onto the second factor. Let $R \subset B^3 \times I$ be a disk with one minimum. That is, R is a properly embedded disk with unknotted boundary in $B^3 \times 1$ so that $h|_R$ is Morse with single index-0 critical point. Let D be a disk in $B^3 \times 1$ bounded by ∂R .




Figure 3.26: A detailed view of a cone singularity in $\mathcal{G}_0$ near the end of band b in a band movie. The two cones have identical neighborhoods, modulo rotation.

Restrict h to $(B^3 \times I) \setminus \nu(R)$. Let $\mathcal{G}_1$ be any singular fibration on $h^{-1}(1)$ with the following properties.

- All but finitely many leaves of $\mathcal{G}_1$ intersect D transversely.
- Where a leaf of $\mathcal{G}_1$ is tangent to D , that leaf locally looks like a minimum or maximum disk or saddle and locally intersects D in a single point.
- No singular points of $\mathcal{G}_1$ meet $\nu(D)$.
- Near $\partial\nu(R)$, the leafs of $\mathcal{G}_1$ intersect D in circles parallel to $\partial\nu(R)$.

By abuse of notation, take a copy of $D := D \setminus \nu(R)$ to live in each $h^{-1}(t)$. Now consult Figure 3.27.

Note $\mathcal{G}_1$ induces a singular foliation $\mathcal{F}$ on D , whose singularities are dots $p_1, \dots, p_k$ and crosses, for $k \geq 1$. Let $p_i \in D$ be a dot singularity in this foliation. Let P_i be the set $\cup \{E \subset D \text{ a disk} \mid p_i \in E, \partial E \text{ in one leaf of } \mathcal{G}_1, p_i \text{ is the unique singularity of}$

$\mathcal{F}|_E\}$. Let $D_i = \bigcup_{E \in P_i} E$. Then either ∂D_i meets a cross singularity of $\mathcal{F}$, or $D_i = D$. Note D_i may not be a disk.

Proposition 3.3. *For some choice of i , D_i is a disk.*

Proof. Suppose D_1 is not a disk. Let E_1 be the interior component of $D \setminus D_1$. E_1 is an open disk, so there is another p_i in E_1 (reorder so $p_2 \in E_1$). Now suppose D_2 is not a disk. Similarly let E_2 be the interior component of $D \setminus D_2$. Then E_2 is an open disk, so there is another p_i in E_2 . Note $E_2 \subset E_1$, so $p_i \neq p_1$ or p_2 (reorder so $p_3 \in E_2$).

We continue inductively. Since there are a finite number of p_i , eventually some D_i must be a disk. $\square$

So without loss of generality, assume D_1 is a disk.

From $t = 1$ to $3/4$, play an interior 0-,1- or 2-,3- stabilization movie in a neighborhood of p_1 (choose according to Fig. 3.28; take the cone to be type II and the dot to be type III). Call the new cone singularity q ; take D to intersect q so that $\mathcal{G}_{3/4}$ induces the same foliation as $\mathcal{G}_1$ on D , but one dot singularity in the foliation $\mathcal{G}_{3/4}|_D$ is at q . From $t = 3/4$ to $t = 1/2$, if $D \neq D_1$ then position the cone along the disk D_1 . (If $D = D_1$, position the cone along a copy of D_1 shrunk slightly to lie in the interior of $h^{-1}(3/4)$.)

Now in $\mathcal{G}_{1/2}$, if $D_1 = D$ then there is a leaf component of $\mathcal{G}_{1/2}$ which is a disk parallel to D . If $D_1 \neq D$, then we can isotope $\mathcal{G}_{1/2}$ so D is disjoint from all singularities of $\mathcal{G}_{1/2}$ and the foliation $\mathcal{G}_{1/2}$ induces on D has $k - 1$ dot singularities.

Repeat this process another $k - 1$ times, from $t = 1/2$ to $t = 1/4$, so that there is a leaf component E of $\mathcal{G}_{1/4}$ which is a disk parallel to D .

Then there is a subset $B \subset h^{-1}(1/2)$ containing E so that $\mathcal{G}_{1/2}$ agrees with $\mathcal{F}_1$ of the minimum movie in B . Play the minimum movie in B from $t = 1/4$ to 0.




Figure 3.27: Left to right, top to bottom: Some leaves of $\mathcal{G}_t$ as t decreases from 1 to 0 in a disk movie. As t decreases, we perform interior stabilizations and position the cones along disks until in a set B containing ∂D in $h^{-1}(1/4)$, $\mathcal{G}_{1/4}$ agrees with the top of the minimum movie. We play the minimum movie from $t = 1/4$ to 0.




Figure 3.28: Leaves near a dot in the singularity $\mathcal{G}_t$ induces on D . The red arrow indicates the positive S^1 direction of $\mathcal{G}_t$. We stabilize, and choose the indices of the stabilization as indicated.

There are exactly k more cone and $k + 2$ more dot singularities in $\mathcal{G}_0$ than in $\mathcal{G}_1$, all of which are types II or III.

In Figure 3.29, we give valid singularity charts for the simple and generalized cancellation movies, and the band and disk movies.

3.7 Proof of Lemma 2.4

Lemma 2.4. Let Z^4 be a compact 4-manifold. Fix a Morse function $h : Z^4 \rightarrow I$. Suppose $\mathcal{F}_t$ is a movie of singular fibrations on Z^4 so that $\mathcal{F}_1$ and $\mathcal{F}_0$ are fibrations (with no singularities) and $\mathcal{F}_t$ is a concatenation of band, disk, and simple or generalized cancellation movies in order (decreasing t).

Then $\mathcal{F}_1 \sqcup \mathcal{F}_0$ extends to a fibration on Z^4 .

Proof. Let $\mathcal{F}_t$ be a singular movie on Z^4 with Morse function $h : Z^4 \rightarrow I$ so that $\mathcal{F}_1$ and $\mathcal{F}_0$ are nonsingular. Assume $\mathcal{F}_t$ is built by composing some number of band, disk, and simple or generalized cancellation movies, in that order (decreasing t).




Figure 3.29: Top left: two singularity charts for the simplified cancellation movie, depending on the indices of the cancelled singularities. Bottom left: two singularity charts for the generalized cancellation movie. Middle: A potential chart for the disk movie, where we included one 0,1- stabilization and one 2,3-stabilization (recall these choices depend on orientations of leaves of $\mathcal{F}_1$ intersecting the disk). Right: two singularity charts for the band movie (recall the types of the cones created in the 1-2- stabilization depends on the orientation of $\mathcal{F}$).

By Theorem 1.11, to conclude that Z^4 is fibered it is sufficient to show that $\mathcal{F}_t$ is valid (perhaps up to reparametrization). In section 3.6, we found singular charts for the band, disk, and generalized cancellation movies.

In particular, the band movie admits a chart in which the two cones at the bottom are type II (opposite indices), and assumes nothing about existing cones and dots.

The disk movie admits a chart in which the cones and dots at the bottom are all type II or III, and assumes nothing about existing cones and dots.

The cancellation movies each admit a chart in which a type II and type III singularity die (the slopes of the corresponding arcs may be positive or negative), and all other arcs are nearly vertical (of arbitrarily-signed slope).




Figure 3.30: A possible valid singularity chart for $\mathcal{F}_t$ when $\mathcal{F}_t$ is a composition of one band movie, one disk movie, and then three cancellation movies. Left: All cancellations are simple. Right: The cancellations are generalized.

Thus, by stacking these charts we obtain a valid singularity chart for $\mathcal{F}_t$. We give two sample singularity charts of $\mathcal{F}_t$ in Figures 3.30 and 3.31. $\square$




Figure 3.31: A possible valid singularity chart for $\mathcal{F}_t$ when $\mathcal{F}_t$ is a composition of two band movies, two disk movie, and then nine cancellation movies. Left: All cancellations are simple. Right: The cancellations are generalized.

Chapter 4

Extending the fibration on the knot complement to a fibration of the disk complement

4.1 Proof of Theorem 1.8

Recall the setup of the ribbon disk D with boundary a fibered knot K . We have:

- $D \subset S^3 \times [1, 3]$.
- For $t_1 > t_2 > \dots > t_n \in (2, 3)$, $D \cap (S^3 \times t_i) =$ a link with a band attached (referred to as a “band,” “ribbon band,” or, “fission band”).
- For $s_1 > \dots > s_{n+1} \in (1, 2)$, $D \cap (S^3 \times s_i)$ is an $(n + 1 - i)$ -component unlink and a disjoint disk (referred to as a minimum disk).
- For all other $u \in (s_{n+1}, 3]$, $D \cap (S^3 \times u)$ is a nonsingular link. For all $u \in [0, s_{n+1})$, $D \cap (S^3 \times u) = \emptyset$.

Theorem 1.8. Suppose the fission bands at $t = t_1, \dots, t_n$ are disjoint and transverse to the fibration on $S^3 \setminus \nu(K)$ (when projected to $S^3 \times 3$). Then the fibration on $S^3 \setminus \nu(K)$ extends to a fibration of handlebodies on $B^4 \setminus \nu(D)$.

Proof. Let $\mathcal{F}_3$ be the fibration on $S^3 \setminus \nu(K) = h^{-1}(3) \setminus \nu(D)$. For $i = 1, \dots, n$, from $t = t_i + \epsilon$ to $t_i - \epsilon$, play a band movie in a neighborhood of b_i . Extend $\mathcal{F}_{t_n - \epsilon}$ vertically to $\mathcal{F}_2 : h^{-1}(2) \setminus \nu(D)$. The singular fibration $\mathcal{F}_2$ has exactly $2n$ cone singularities and zero dot singularities.

Let $d_1, \dots, d_{n+1}$ be disjoint disks in $S^3 \times 2$ with d_i corresponding to the minimum at s_i . For $i = 1, \dots, n$, from $t = s_i + \epsilon$ to $s_i - \epsilon$, play a disk movie near each d_i . Extend $\mathcal{F}_t$ vertically to $\mathcal{F}_{s_n - \epsilon}$ (that is, just past the penultimate minimum). The singular fibration $\mathcal{F}_{s_n - \epsilon}$ has an equal number of cone and dot singularities. (Say $\mathcal{F}_{s_n - \epsilon}$ has r cone singularities.)

Finally, we play r generalized cancellation movies to extend $\mathcal{F}_t$ to a movie on $h^{-1}([s_{n+1} + \epsilon, 3]) \setminus \nu(D)$ so that $\mathcal{F}_{s_{n+1} + \epsilon}$ has no singularities.

By Lemma 2.4, $(B^4 \setminus \nu(D)) \cap h^{-1}([s_{n+1} + \epsilon, 3])$ admits a fibration $\mathcal{F}$. Note $(B^4 \setminus \nu(D)) \cap h^{-1}([0, s_{n+1}]) \cong (B^4 \setminus (\text{trivial disk})) \cong S^1 \times B^3$. Glue $\mathcal{F}$ to the fibration of $S^1 \times B^3$ by 3-balls to obtain a fibration $\mathcal{G}$ of $B^4 \setminus \nu(D)$.

It follows immediately from [16, Theorem 1.1] that the fibers of $\mathcal{G}$ are handlebodies, completing the proof of Theorem 1.8. $\square$

Remark 4.1. It is essential that D is a ribbon disk and that the fission bands defining D are transverse to the fibration. These conditions ensure that after extending the movie of fibrations below the heights of all bands and all but one minimum disk, there are an equal number of cone and dot singularities in the cross-section $\{t = s_n - \epsilon\}$ (above the bottom minimum disk).

In section 4.3, we show explicitly that the fibers of $\mathcal{G}$ are handlebodies. In chapter 5, we show how to obtain 2-handle attaching circles for a handlebody fiber H of

$\mathcal{G}$ on $\partial H \cap h^{-1}(3) = (\text{a fiber for } K)$. This explicitly describes the embedding of H into $B^4 \setminus \nu(D)$ (up to isotopy rel boundary).

4.2 Remark on the generalized cancellation movies

In the proof of Theorem 1.8, the generalized cancellation movies played from $t = s_n - \epsilon$ to $s_n + \epsilon$ may actually be taken to be simple cancellation movies. This is not strictly necessary for Theorem 1.8, but is necessary to explicitly understand the embedding of a handlebody fiber of D into $B^4 \setminus \nu(D)$, which we will discuss in chapter 5. We first give a more general statement.

Lemma 4.2. *Let V be a connected 3-manifold with torus boundary, and $\mathcal{F}_1$ be a singular fibration on V with r cone and r dot singularities, which are all type II or III. Assume that $\mathcal{F}_1^{-1}(\theta) \cap \partial V$ is a connected simple closed curve for each θ . Then by playing only simple cancellation movies, $\mathcal{F}_1$ can be extended to a valid movie of singular fibrations $\mathcal{F}_t$ on $V \times I$ so that $\mathcal{F}_0$ is nonsingular.*

Proof. Note that $\mathcal{F}_1$ can be extended to a movie $\mathcal{F}_t$ in which $\mathcal{F}_0$ is nonsingular via *generalized* cancellation movies. Therefore, $V = V \times 0$ admits a fibration over S^1 . Then V is an irreducible 3-manifold (with boundary), and in particular every 2-sphere embedded in V bounds a 3-ball.

Proposition 4.3. *If $\mathcal{F} := \mathcal{F}_1$ is nonsingular, then it has a dot singularity p so that when flowing from p outward perpendicular to the leaves of $\mathcal{F}$, the first singular leaf component has a cone q so that the following are true:*

- *With respect to the circular Morse function $\mathcal{F}$, p and q are critical points of index $(0,1)$ or $(3,2)$.*
- *Suppose $\mathcal{F}^{-1}(\theta) \cap \nu(q)$ is a two-sheeted hyperboloid. The two sheets are not in the same component of $\mathcal{F}^{-1}(\theta)$.*

Proof. Suppose $\mathcal{F}$ has k_i critical points of index i . Then $0 = \chi(V) = k_0 - k_1 + k_2 - k_3$. But recall there are an equal number of dot and cone singularities in V , so $k_0 + k_3 = k_1 + k_2$. Therefore, $k_0 = k_1$ and $k_2 = k_3$.

Let p be a dot singularity of $\mathcal{F}$, and reorient $\mathcal{F}$ if necessary so that p is an index-0 critical point of $\mathcal{F}$. Flow from p in the positive S^1 direction until finding a singular leaf (with cone point q). This must happen eventually, since $\mathcal{F}$ is defined on a manifold with boundary.

Claim. If q is a cone of index 1, then q satisfies the second condition of the proposition.

Proof. We argue that every index-1 cone satisfies the second condition.

Recall $k_0 = k_1$. Fix a nonsingular leaf L of $\mathcal{F}$ which meets ∂V . Reparametrize $\mathcal{F}$ in a neighborhood of the index-0 points so that $\mathcal{F}^{-1}(0) = L \cup \{\text{the } k_0 \text{ index-0 critical points}\}$ and the cone singularities all lie in distinct leaves of $\mathcal{F}$. Let $V_\theta = \cup_{0 \leq x \leq \theta} \mathcal{F}^{-1}(x)$. Suppose V_θ has n components for $\theta_0 - \epsilon < \theta < \theta_0$ while V_θ has $n - 1$ components for $\theta_0 \leq \theta < \theta_0 + \epsilon$, for some θ_0 . Then $\mathcal{F}^{-1}(\theta_0)$ must contain a cone satisfying the proposition. See Figure 4.1. (Recall each $\mathcal{F}^{-1}(\theta)$ meets ∂V in one curve.)

Note $V_0 = \mathcal{F}^{-1}(0)$ has at least $k_0 + 1$ components, while $V_{2\pi} = V$ has one component. Therefore, the number of components must decrease at least k_0 times. But there are $k_1 = k_0$ cones, so every index-1 cone must satisfy the second condition of proposition. $\square$

Now assume q is an index-2 critical point. See Figure 4.2. There is a leaf component of $\mathcal{F}$ near q which is sphere separating V into two pieces, one of which is a ball B not containing p . There is some dot p' inside B . Repeat the argument starting from the dot p' instead of B . If p' does not satisfy Proposition 4.3, then we find another spherical leaf component of $\mathcal{F}$ inside B , bounding a ball in $B' \subset B$ that does




Figure 4.1: Schematic of V_θ for increasing θ . Left: V_θ for $\theta > 0$ small. Middle: The number of components of V_θ decreases. $\mathcal{F}^{-1}(\theta)$ contains a cone satisfying the second condition of the proposition. Right: We draw two cones. One is index-2, one is index-1 but does not satisfy the other condition. The number of components of V_θ does not decrease.

not contain p' . Then there is a dot point p'' inside B' . We continue inductively; since there are finitely many critical points in $\mathcal{F}$, eventually we must find a dot point as in Proposition 4.3. $\square$

Let p, q satisfy Proposition 4.3. Let $\eta : [0, 1]$ be an arc from q to p intersecting the leaves of $\mathcal{F}_1$ transversely. Proposition 4.3 exactly ensures that we may play a simple cancellation movie around η to extend $\mathcal{F}_1$ to a valid movie on $t \in [1 - \epsilon, 1]$ so that $\mathcal{F}_{1-\epsilon}$ has $r - 1$ cone and $r - 1$ dot singularities. Proceed inductively to extend $\mathcal{F}_t$ to a valid movie on $t \in [0, 1]$ so that $\mathcal{F}_0$ is nonsingular. $\square$

In the setting of Theorem 1.8, $\mathcal{F}_{s_n - \epsilon}$ is a singular fibration of a solid torus satisfying the hypotheses of Lemma 4.2. Therefore, we may take all the cancellation movies played from $t = s_n - \epsilon$ to $s_{n+1} + \epsilon$ to be simple.




Figure 4.2: Starting from dot p and flowing perpendicular to the singular fibration $\mathcal{F}$, the next singular leaf we find has a cone q corresponding to an index-2 critical point. A nearby leaf component is a sphere bounding ball B containing dot p' . Starting from p' and flowing perpendicular to the leaves of $\mathcal{F}$, again the first cone corresponds to a critical point of index-2. A nearby leaf component is a sphere bounding ball $B' \subset B$ containing dot p'' .

4.3 Understanding the fibers of $B^4 \setminus \nu(D)$

In this section, we explicitly show that the fibers of $\mathcal{G}$ are handlebodies, via the construction of Theorem 1.8 and Theorem 1.11.

Let H be the fiber of $B^4 \setminus \nu(D)$ obtained from Theorem 1.8. Then $F = H \cap \partial B^4$ is a leaf in the fibration of $S^3 \setminus \nu(K)$. Explicitly, $H \cong F \times I \cup \{1\text{-}, 2\text{-}, 3\text{-handles}\}$, where the 1-handles correspond to the type I cones which appear and vanish during a band movie.

If a minimum disk d_i intersects the interior of band b_j (when both are projected to the same height), then when a disk movie is played near d_i , we obtain 2-handles in H geometrically cancelling all the 1-handles corresponding to the band movie for band b_j . See Figure 4.3. We must now only be careful that a different minimum disk cancel the 1-handles for each band.




Figure 4.3: Rectangular sheet: part of the minimum disk d_i which intersects a core of the band b_j . Surface: A leaf L of $\mathcal{F}_t$ for $t = t_i - \epsilon$. Some curves in $L \cap d_i$ run along the cores of the 1-handles corresponding to b_j (attached to the 3-dimensional fiber H). These curves are all distinct, since they locally have the same orientation in d_i (Here, we are using the fact that b_j is transverse to the fibration $\mathcal{F}_t$). Therefore, 2-handles attached to H corresponding to the Type II cones arising during the disk movie on d_i will geometrically cancel the 1-handles corresponding to the type I cones arising during the band movie on b_j .

Let $d_1, \dots, d_{n+1}$ be the minimum disks of D (projected into $h^{-1}(1)$).

Lemma 4.4. *Up to reordering the minimum disks $\{d_i\}$, we may assume $d_i \cap \mathring{b}_i \neq \emptyset$.*

Proof. Let Γ be a bipartite graph with $2n + 1$ vertices in $V \cup W = \{v_1, \dots, v_{n+1}\} \cup \{w_1, \dots, w_n\}$. Say v_i and w_j are adjacent if $d_i \cap \mathring{b}_j \neq \emptyset$. If $\partial d_i \cap \partial b_j \neq \emptyset$ but $d_i \cap \mathring{b}_j = \emptyset$, then we may isotope D to cancel d_i and b_j (without moving the other bands). Assume we have cancelled all such pairs.

Let $X \subset W$ be a set of $k \leq n$ vertices. Let $N_\Gamma(X)$ denote the subset of V which is adjacent to X . Suppose $|N_\Gamma(X)| < k$. Then some band b_i with $w_i \in X$ is a fusion band, rather than fission, yielding a contradiction. Thus, by Hall's Marriage Theorem, there exists a perfect matching between W and V . That is, we can relabel the disks d_i (and corresponding vertices w_i) so that $d_i \cap \mathring{b}_i \neq \emptyset$ (i.e. v_i is adjacent to w_i). □

By Proposition 4.4, a collection of 2-handles in the fiber H of $\mathcal{G}$ on $B^4 \setminus \nu(D)$ (resulting from the minimum disk d_i) geometrically cancel all the 1-handles corresponding to the band b_i . Therefore, $H \cong (\partial H \times I) \cup \{2\text{-,}3\text{-handles}\}$, so H is a handlebody. $\square$

4.4 Proof of Theorem 1.10

Theorem 1.10. Let K be a fibered knot in S^3 . Let $J \subset S^3$ be a knot so that there is a *ribbon concordance* from K to J (i.e. there is an annulus A properly embedded in $S^3 \times I$ with boundary $(K \times 1) \sqcup (\bar{J} \times 0)$ so that $\text{proj}_I|_A$ is Morse with no local maxima). Say the index-1 critical points of A correspond to fission bands attached to K . If the bands can be isotoped to be transverse to the fibration on $S^3 \setminus \nu(K)$, then $(S^3 \times I) \setminus \nu(A)$ is fibered by compression bodies.

Proof. We proceed as in Theorem 1.8. Let A be an annulus in $S^3 \times I$ with boundary $K \times 1 \sqcup \bar{J} \times 0$, with no local maxima with respect to projection on I . Say the local minima of A occur in $S^3 \times s_i$ for $s_1 > s_2 > \dots > s_n \in [2/7, 3/7]$ and the index-1 critical points in $S^3 \times t_i$ for $t_1 > t_2 > \dots > t_n \in [5/7, 6/7]$. By hypothesis, we assume that bands corresponding to the index-1 critical points are transverse to the fibration $\mathcal{F}_1$ on $(S^3 \times 1) \setminus \nu(K)$ (when projected to $S^3 \times 1$).

We build a movie of singular fibrations $\mathcal{F}_t$ on $(S^3 \times I) \setminus \nu(A)$ as in Theorem 1.8. Extend $\mathcal{F}_t$ to $(S^3 \times [4/7, 1]) \setminus \nu(A)$ by playing a band movie near each band of A . Then extend to $(S^3 \times [1/7, 1]) \setminus \nu(A)$ by attaching playing a disk movie near each minimum disk of A . There are an equal number of cone and dot critical points in $\mathcal{F}_{1/6}$, so we may play generalized cancellation movies to extend $\mathcal{F}_t$ to $(S^3 \times I) \setminus \nu(A)$ with $\mathcal{F}_0$ a nonsingular fibration. (In fact, these generalized cancellations could be taken to be simple by Lemma 4.2.) Thus, J is fibered.

By Lemma 2.4, $(S^3 \times I) \setminus \nu(A)$ admits a fibration $\mathcal{G}$ with fiber H . Via the argument of section 4.3, $H \cong ([H \cap (S^3 \times 1)] \times I) \cup \{2\text{- and } 3\text{-handles}\}$. Therefore, H is a compression body from the fiber for K to the fiber for J .

Note $H \cong (\overset{\circ}{\Sigma}_{g(K)} \times I) \cup \{g(K) - g(J) \text{ 2-handles}\} \cong \natural_{g(K)+g(J)} S^1 \times D^2$.

□

Chapter 5

Conclusion: explicitly finding the fiber of $B^4 \setminus \nu(D)$

Let F be a fiber for fibered knot K . Let D be a ribbon disk for D so that fission bands b_i for D are transverse to the fibration of $S^3 \setminus \nu(K)$. By Theorem 1.8, the fibration extends to a fibration $\mathcal{G}$ on $B^4 \setminus \nu(D)$ with fiber H , where $\partial H = F \cup D$.

In the proof of Theorem 1.8, we constructed H by building a valid movie $\mathcal{F}_t, t \in [s_{n+1} + \epsilon, 3]$ on $B^4 \setminus \nu(D)$ so that $H = \cup_t \mathcal{F}_t^{-1}(\theta) \cup (3\text{-handle})$ for some θ_0 . The 2-handles of H correspond to the type II cones and type II_B of $\mathcal{F}_t$ intersecting the vertical $\theta = \theta_0$ line in a valid chart $\mathcal{C}$ for $\mathcal{F}_t$.

Recall $\mathcal{F}_t$ may be built from band, disk, and simple (see section 4.2) cancellation movies. Say that from $t = 3$ to 2, $\mathcal{F}_t$ is built from band movies; from $t = 2$ to 1, $\mathcal{F}_t$ is built from disk movies; from $t = 1$ to $s_{n+1} + \epsilon$, $\mathcal{F}_t$ is built from simple cancellation movies. In $\mathcal{C}$, every type II cone arises during a band movie or a disk movie. The type II cones arising in a band movie correspond to arcs which are nearly vertical until $t = 1$. From $t = 1$ to $t = s_{n+1} + \epsilon$, during simple cancellation movies all cones correspond to arcs which are nearly vertical. Therefore, we may choose θ_0 so that the vertical $\theta = \theta_0$ line intersects type II arcs only in $1 < t < 2$, does not intersect




Figure 5.1: We draw potential valid singularity charts for $\mathcal{F}_t$, $t \in [s_{n+1}, 3]$. By Lemma 4.2, we may assume that the cancellations of $\mathcal{F}_t$ are all simple, as on the left. Then all 2-handles of H arise from positioning a cone or playing a minimum movie during a disk movie. Therefore, the 2-handles are attached exactly along the intersections of $\mathcal{F}_t^{-1}(\theta)$ with a minimum disk, projected to the same height.

the arcs of type II cone singularities arising from band movies, and does not meet any type II_B singularities. (Perhaps after reparametrizing $\mathcal{F}_t$ so that type II_B and II boundary arcs are short.) See Figure 5.1 (left).

To recap, this means that every type II or II_B singularity meeting the line $\theta = \theta_0$ in $\mathcal{C}$ arises during a disk movie. In $H \cap h^{-1}(t)$, as t decreases, passing through these singularities corresponds to compressing $H \cap h^{-1}(t)$ along a closed curve of intersection with a minimum disk of D . We conclude that intersections of the minima of D (projected to S^3) with $H \cap h^{-1}(3) = F$ give attaching circles of the 2-handles of H .

(This decomposition is not minimal if the line $\theta = \theta_0$ meets type III arcs in $\mathcal{C}$, but we may choose a set of linearly independent curves to give 2-handles if we wish.)

Therefore, to find the 2-handle attaching curves on $F \cup D$ defining H , we do the following:

- Find a disjoint collection of disks $d_1 \sqcup \cdots \sqcup d_n$ in S^3 bounded by K resolved along the b_i .
- Isotope the d_i so that near the interior of $\partial d_i \cap \partial F$, d_i and F coincide.
- Draw a (maximal linearly independent) collection of closed curves $d_i \cap F$ on F .

These are 2-handle attaching curves on $F \cup D$ which yield H .

We could equivalently state this process as:

- Project D to S^3 to find an immersed ribbon disk D' with boundary K .
- Isotope D' so that near $\partial D' = \partial F$, F coincides with the sheet of D' containing the boundary.
- Draw a (maximal linearly independent) collection of closed curves in $D' \cap F$ on F . These are 2-handle attaching curves on $F \cup D$ which yield H .

In Figure 5.2, we give explicitly a fibered ribbon disk and use the above procedure to find attaching circles for 2-handles in the handlebody fiber. In Figure 5.3 we repeat this process for two fibered ribbon disks with the same boundary to find the fiber of a fibered 2-knot in S^4 .




Figure 5.2: Left: The fibered knot $K = 8_{20}$ (i.e. the pretzel knot $P(2, -3, 3)$). We draw a fission band defining a ribbon disk D with boundary K and shade a genus-2 fiber surface for K . On the fiber, we draw two attaching circles defining a genus-2 handlebody fiber of $B^4 \setminus \nu(D)$. Right: Two disjoint disks bounded by K resolved along the fission band. These disks intersect the fiber in arcs (near the band) and two curves, which are the attaching circles drawn on the left.




Figure 5.3: Left: A diagram for the fibered ribbon knot $K = 11n_{74}$ (i.e. the pretzel knot $P(3, -3, 3, -2)$). Each of the pictured bands specifies a 2-minimum ribbon disk D_i for K . (These examples of disks come from [20], who produce many examples of doubly slice knots.) Using the construction of Theorem 1.8, we find attaching circles for the handlebody fiber for each disk. Middle: The fiber F for K consists of two annuli glued along three half-twisted bands. The two sets of circles specify the handlebody fiber for the disks given by the bands. Right: On $F \cup D$, the attaching circles specify a Heegaard splitting of S^3 . Thus, $D_1 \cup \overline{D_2} \subset S^4$ is fibered by 3-balls (so $D_1 \cup \overline{D_2}$ is the unknotted sphere).

Chapter 6

More examples

It remains open whether any/every ribbon disk for a fibered knot K in S^3 can be defined by bands transverse to the fibration of $S^3 \setminus \nu(K)$. However, for specific examples of K we can check that this condition is satisfied by some ribbon disk D .

In particular, we find fibered ribbon disks for small fibered ribbon knots.

Proposition 6.1. *Let K be a prime fibered ribbon knot with crossing number at most 12. Then K bounds a ribbon disk $D \subset B^4$ so that $S^4 \setminus \nu(D)$ is fibered by handlebodies.*

Proof. By Corollary 1.14, if K bounds a disk D with two minima, then the fibration on $S^3 \setminus \nu(K)$ extends to a fibration of $B^4 \setminus \nu(D)$. In particular, every prime ribbon knot of at most 10 crossings bounds a 2-minimum disk [13], as do most prime fibered knots of 11 and 12 crossings (observed through e.g. symmetric union diagrams [15]). Using the KnotInfo database [4] to list fibered slice knots, and symmetric union diagrams of these knots due to Lamm [15], we explicitly produce a 3-minimum disk D for each 11- or 12-crossing prime fibered ribbon knot which does not bound an obvious-to-the-author 2-minimum disk. The bands for each disk either lie in a fiber for the knot $K \subset S^3$, or run transverse to the fibration. See Figure 6.1. Transversality of a band can be seen by observing the band core is given by taking an arc in a fiber and adding twists (of one sign) around the binding (knot).

In the first draft of this paper, the author was not aware of the following examples of slice disks due to Brendan Owens and Frank Swenton (as of this writing, they have not yet released a paper including these disks, but Brendan Owens kindly let this author know of their examples). They have constructed (in particular) examples of 2-minimum disks bounded by each of the knots in Figure 6.1 except for $12a_{990}$. (These are each potentially isotopic to the 3-minimum examples.) We retain the 3-minimum examples as illustrations of how one can verify the transversality condition of Theorem 1.8. In Figure 6.2, we provide the 2-minimum disks of Owens and Swenton. □

Question 6.2. *Does $12a_{990}$ bound a 2-minimum ribbon disk?*

Note here we defined “small” knots to be prime knots with fewer than 13 crossings. One might reasonably consider “small” knots to be 2-bridge knots. In Lisca’s [19] classification of ribbon 2-bridge knots, he produces an explicit ribbon disk for every ribbon 2-bridge knot. Most of these disks have two minima, except for the examples in Figure 6.3 (left). By drawing the ribbon knots as symmetric unions (using diagrams of Cristoph Lamm¹), we find ribbon disks with two minima (Fig. 6.3, right). Therefore, by Corollary 1.14, every fibered ribbon 2-bridge knot bounds a fibered ribbon disk.

However, we note that in this setting, the 2-bridge notion of “small” is actually less interesting than low crossing-number. In fact, there are finitely many fibered ribbon 2-bridge knots. We are not aware of this fact in the literature, so we include this as an interesting remark.

Remark 6.3. Up to mirroring, there are exactly five fibered ribbon 2-bridge knots: $8_9, 9_{27}, 10_{42}, 11a_{96}, 12a_{477}$.

¹This preprint is available online as C. Lamm, *Symmetric union presentations for 2-bridge ribbon knots*, arXiv:math/0602395 [math.GT], Feb. 2006




Figure 6.1: These prime fibered ribbon knots did not obviously (to the author) bound 2-minimum disks in B^4 . We produce here a 3-minimum ribbon disk for each knot. The bands for $11a_{164}$, $11a_{326}$, $12a_{458}$, $12a_{473}$, $12a_{887}$, $12a_{1225}$ each lie in the illustrated fiber. Knots $12a_{427}$, $12a_{101}$, $12a_{1105}$ are drawn with one band in the fiber and one transverse to the fibration. Both bands for $12a_{990}$ run transverse to the fibration.




Figure 6.2: A band diagram of a 2-minimum disk for each knot of Fig. 6.1 except $12a_{990}$, constructed by B. Owens and F. Swenton.




Figure 6.3: Left: bands defining a ribbon disk for $K \subset S^3$, constructed by Lisca [19]. Right: A 2-minimum ribbon disk with the same boundary knot. Note here that the integers $\pm 2a, \pm 2b$ refer to half-twists. The knot K is fibered if and only if $|a| = |b| = 1$.

Proof. By Casson and Gordon [3], every ribbon 2-bridge knot (up to mirroring) has a continued fraction expansion in one of the following three families:

- Family 1: $[c_1, -c_2, c_3, -c_4, \dots, c_{2n+1}, -1, -c_{2n+1}, \dots, c_4, -c_3, c_2, -c_1]$ where each $c_i > 0$,
- Family 2: $[2a, 2, 2b, -2, -2a, -2b]$, $a, b \neq 0$,
- Family 3: $[2a, 2, 2b, -2a, -2, -2b]$, $a, b \neq 0$.

As this is somewhat of a side note from the main thrust of this paper, we omit exposition of continued fraction expansion notation for 2-bridge knots. For those interested, we recommend section 2 of [12].

Proposition 6.4. *A knot K in Family 1 is never fibered.*

Proof. Let $[d_1, \dots, d_m]$ be a *strict* continued fraction for $[-c_{2n+1}, \dots, c_4, -c_3, c_2, -c_1]$, as in [12] (i.e. all d_i are nonzero, d_{2i+1} is even and if $|d_{2i+1}| = 2$ then $d_{2i+1}d_{2i+2} < 0$). Then K has continued fraction expansion $[-d_m, \dots, -d_1, -1, d_1, \dots, d_m]$. For odd m , this fraction gives a 2-component link, so m must be even. We draw a minimum-genus Seifert surface F for K in Figure 6.4. From F , we can deplumb an annulus with $d_1 - \text{sign}(d_2) + 1$ half-twists and an annulus with $-(d_1 - \text{sign}(d_2)) + 1$ half-twists. These cannot both be Hopf bands, so F is not a fiber. $\square$

In Figure 6.5, we draw a minimum-genus Seifert surface for a knot in Family 2 or 3. In either case, the surface decomposes as a Murasugi sum of two Hopf bands and 4 annuli with $2a, 2b, -2a, -2b$ half-twists, respectively. Therefore, the knot is fibered if and only if $|a| = |b| = 1$.

The Family 2 knot with $a = 1, b = -1$, and the Family 3 knots with $\{a, b\} = \{1, -1\}$ are all 9_{27} . The Family 2 knot with $a = 1, b = 1$ and the Family 3 knot with $a = 1, b = 1$ are both 8_9 . There are no other repeats, so we are left




Figure 6.4: A minimum-genus Seifert surface for 2-bridge ribbon knots in Family 1. From this surface, we can deplumb annuli with $d_1 - \text{sign}(d_2) + 1$ and $-d_1 + \text{sign}(d_2) + 1$ half-twists. These cannot both be Hopf bands, so the surface is not a fiber (and hence the knot is not a fibered knot).




Figure 6.5: Minimum-genus surfaces for knots in Families 2 and 3. These are fibers if and only if $|a| = |b| = 1$, so a knot in Family 2 or 3 is fibered if and only if $|a| = |b| = 1$.

with five distinct 2-bridge knots. Using KnotFinder [4], we identify these knots as $8_9, 9_{27}, 10_{42}, 11a_{96}, 12a_{477}$. □

Chapter 7

More questions and final comments

Recall Question 1.5, the main motivation for this paper.

Question 1.5. Let D be a homotopy-ribbon disk in B^4 so that $\partial D \subset S^3$ is a fibered knot. Is $B^4 \setminus \nu(D)$ fibered?

This naturally leads to the (potentially) easier question:

Question 7.1. *Let K be a fibered homotopy-ribbon knot in S^3 . Does K bound a fibered homotopy-ribbon disk in B^4 ?*

We remark that if the answer to Question 7.1 is “no” for any knot K , then the relative smooth 4-dimensional Poincaré conjecture is false.

Proof. Assume K is fibered and homotopy-ribbon in S^3 . By Casson and Gordon [2], K bounds a disk D fibered by handlebodies in a homotopy 4-ball V . By Larson and Meier [16], if $V \cong B^4$ then D is strongly homotopy-ribbon. $\square$

In Remark 1.6, we claimed that the converse of Larson and Meier’s theorem holds even when “strongly homotopy-ribbon” is relaxed to “homotopy-ribbon.” That is, if a disk D in B^4 is homotopy-ribbon and fibered, then the fibers are handlebodies. This follows easily from the proof of [16] and a paper by Cochran [5] cited therein, but

we sketch the proof here as a matter of interest. This proof closely follows arguments in both of these papers; one should refer to [16] or [5] for more detail and exposition on disk-knots in B^4 or ribbon 2-knots in S^4 , respectively.

Proof. We make use of the following lemma of Cochran: Let M be a closed 3-manifold. Then the natural map $i^\# : M \rightarrow K(\pi_1(M), 1)$ induces the zero map $i_* : H_3(M) \rightarrow H_3(\pi_1(M))$ if and only if $M = \#_h S^1 \times S^2$ for some $h \geq 0$.

Let D be a fibered homotopy-ribbon disk in B^4 , with fiber H . Then $K = \partial D$ is a fibered knot, with fiber F .

Let $W = B^4 \setminus \nu(D)$ and $\widetilde{W}$ be the infinite cyclic cover of W . Then $\widetilde{W} \cong H \times \mathbb{R}$, $\partial \widetilde{W} \cong F \times \mathbb{R}$. Thus, $\pi_1(H) \cong [\pi_1(W), \pi_1(W)]$ and $\pi_1(F) \cong [\pi_1(S_0^3(K)), \pi_1(S_0^3(K))]$. Since D is homotopy-ribbon, the inclusion map $\pi_1(S^3 \setminus K) \rightarrow \pi_1(W)$ is surjective. Therefore, the inclusion $\pi_1(F) \rightarrow \pi_1(H)$ is surjective.

Let G be the 2-sphere $D \cup \overline{D}$ in $S^4 = B^4 \cup \overline{B^4}$. Then $S^4 \setminus \nu(G)$ is fibered, with fiber $M' = H \cup_{F=\overline{F}} \overline{H}$. Let $M = M' \cup B^3$ be a closed 3-manifold. By Seifert van-Kampen, the inclusion $H \hookrightarrow M$ induces an isomorphism $\pi_1(H) \cong \pi_1(M)$.

Let X be obtained by surgering S^4 along G (i.e. deleting $\nu(G)$ and gluing in $S^1 \times B^3$). Let $A = (B^4 \times I) \setminus (D \times I)$ be a 5-manifold with $\partial A \cong X$. The inclusion $\widetilde{X} \hookrightarrow \widetilde{A}$ induces an isomorphism $\pi_1(\widetilde{X}) \cong \pi_1(\widetilde{A})$, since $\pi_1(\widetilde{A}) = \pi_1(\widetilde{B^4 \setminus D}) \cong \pi_1(H) \cong \pi_1(M) \cong \pi_1(\widetilde{X})$. We obtain the following commutative diagram (left), which induces a commutative map of spaces (right).

$$\begin{array}{ccc}
 \pi_1(\widetilde{A}) & & \widetilde{A} \\
 i_* \uparrow & \searrow & \uparrow i \\
 \pi_1(\widetilde{X}) & \xrightarrow{\text{id}} [\pi_1(X), \pi_1(X)] & \widetilde{X} \xrightarrow{i^\#} K([\pi_1(X), \pi_1(X)], 1)
 \end{array}$$

Since $H_3(\widetilde{A}) = H_3(\widetilde{B^4 \setminus D}) \cong H_3(H \times \mathbb{R}) \cong H_3(H) = 0$ and $i_* : H_3(\widetilde{X}) \rightarrow H_3(\pi_1(X), \pi_1(X))$ factors through $H_3(\widetilde{A})$, i_* is the zero map. But $\widetilde{X} \cong \mathbb{R} \times \widehat{M}$ and

$[\pi_1(X), \pi_1(X)] \cong \pi_1(M)$, so $i_* : H_3(M) \rightarrow H_3(\pi_1(M))$ is the zero map. Therefore, $M \cong \#_g S^1 \times S^2$ for some g . Since $\pi_1(F) \cong \pi_1(H)$, g is the genus of F and H is a genus- g handlebody.

□

The following more restrictive version of Question 7.1 may be easier to answer.

Question 7.2. *Let K be a fibered ribbon knot in S^3 . Does K bound a fibered ribbon disk in B^4 ?*

In chapter 6, we gave an affirmative answer to Question 7.2 when K is prime with fewer than 13 crossings.

Let K be a ribbon knot. Let S be a minimum-genus Seifert surface for K . Let D be a ribbon disk for K , and project D to S^3 to find an immersed ribbon disk D' in S^3 bounded by K . Isotope S so that S agrees with one sheet of D' near K . Without K and the ribbon disk D obeying the hypotheses of Theorem 1.8, it is not true that $\{D' \cap S\}$ must contain $g(S)$ linearly independent curves in $H_1(S; \mathbb{Z})$ (such a set of curves is called a “ribbon derivative” for K on S). Cochran and Davis [6] constructed genus-1 ribbon knots whose genus-1 Seifert surfaces do not admit ribbon derivatives. Of course, these examples are not fibered (as the only genus-1 fibered knots are the two trefoils and the figure eight, none of which are slice).

Question 7.3. *If K is fibered (and hence S is a fiber), then must $S \cap D'$ include $g(S)$ simple closed curves which are linearly independent in $H_1(S; \mathbb{Z})$?*

We used the transversality of b_i to the fibration on $S^3 \setminus \nu(K)$ to ensure the 1-handles in the relative handlebody decomposition of H are all geometrically cancelled. Without the transversality condition, we might amend the band movie to find a 3-manifold bounded by D (although not a fibration of $B^4 \setminus \nu(D)$). However, we have not shown that a general ribbon disk D bounds a *handlebody* in B^4 .

Question 7.4. *Let $D \subset B^4$ be a ribbon disk for a knot K . What conditions on D and K imply that D bounds a handlebody in B^4 ?*

In the case that ∂D is the unknot, proving that D bounds 3-ball as in Question 7.4 would positively answer the following conjecture¹.

Suzuki's unknotting conjecture [27]. Let G be a 2-knot in S^4 with unknotted equator. Then G is smoothly unknotted.

We remark that if a 2-knot G has unknotted equator, then $\pi_1(S^4 \setminus G) \cong \mathbb{Z}$, so Suzuki's unknotting conjecture is a subcase of the following more general conjecture.

Smooth 4-dimensional unknotting conjecture (see [14], Problem 1.55(a)). Let G be a 2-knot in S^4 with $\pi_1(S^4 \setminus G) \cong \mathbb{Z}$. Then G is smoothly unknotted.

The 3-dimensional unknotting conjecture (Dehn's lemma) was proved by Papakyriakopoulos [23]. For $n > 4$, the n -dimensional unknotting conjecture was proved in the topological case by Stallings [26] and later in the smooth case by Levine [17] [18].

By work of Freedman and Quinn [9], if $\pi_1(S^4 \setminus G) \cong \mathbb{Z}$, then G is topologically unknotted (i.e. G bounds a 3-ball topologically embedded into S^4). Whether G is smoothly unknotted remains an open question.

¹The author strongly recommends the reference [27] to anyone interested in 2-dimensional knot theory.

Bibliography

- [1] Gerhard Burde, Heiner Zieschang, and Michael Heusener. *Knots*, volume 5 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, extended edition, 2014.
- [2] A. J. Casson and C. McA. Gordon. A loop theorem for duality spaces and fibred ribbon knots. *Invent. Math.*, 74(1):119–137, 1983.
- [3] A. J. Casson and C. McA. Gordon. Cobordism of classical knots. In *À la recherche de la topologie perdue*, volume 62 of *Progr. Math.*, pages 181–199. Birkhäuser Boston, Boston, MA, 1986. With an appendix by P. M. Gilmer.
- [4] J.C. Cha and C. Livingston. Knotinfo: Table of knot invariants. <http://www.indiana.edu/~knotinfo>. Accessed 9 August 2018.
- [5] Tim Cochran. Ribbon knots in S^4 . *J. London Math. Soc. (2)*, 28(3):563–576, 1983.
- [6] Tim D. Cochran and Christopher William Davis. Counterexamples to Kauffman’s conjectures on slice knots. *Adv. Math.*, 274:263–284, 2015.
- [7] R. H. Fox. A quick trip through knot theory. In *Topology of 3-manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961)*, pages 120–167. Prentice-Hall, Englewood Cliffs, N.J., 1962.
- [8] R. H. Fox. Some problems in knot theory. In *Topology of 3-manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961)*, pages 168–176. Prentice-Hall, Englewood Cliffs, N.J., 1962.
- [9] Michael H. Freedman and Frank Quinn. *Topology of 4-manifolds*, volume 39 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1990.
- [10] David Gabai. Detecting fibred links in S^3 . *Comment. Math. Helv.*, 61(4):519–555, 1986.
- [11] Jonathan Hillman. *Algebraic invariants of links*, volume 52 of *Series on Knots and Everything*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, second edition, 2012.
- [12] Mikami Hirasawa and Kunio Murasugi. Genera and fibredness of Montesinos knots. *Pacific J. Math.*, 225(1):53–83, 2006.

- [13] Akio Kawauchi. *A survey of knot theory*. Birkhäuser Verlag, Basel, 1996. Translated and revised from the 1990 Japanese original by the author.
- [14] Rob Kirby, editor. *Problems in low-dimensional topology*. Citeseer, 1995.
- [15] Cristoph Lamm. The search for non-symmetric ribbon knots. To appear in *Exp. Math.*
- [16] Kyle Larson and Jeffrey Meier. Fibered ribbon disks. *J. Knot Theory Ramifications*, 24(14):1550066, 22, 2015.
- [17] J. Levine. Unknotting spheres in codimension two. *Topology*, 4:9–16, 1965.
- [18] J. Levine. An algebraic classification of some knots of codimension two. *Comment. Math. Helv.*, 45:185–198, 1970.
- [19] Paolo Lisca. Lens spaces, rational balls and the ribbon conjecture. *Geom. Topol.*, 11:429–472, 2007.
- [20] Charles Livingston and Jeffrey Meier. Doubly slice knots with low crossing number. *New York J. Math.*, 21:1007–1026, 2015.
- [21] John Milnor. *Singular points of complex hypersurfaces*. Annals of Mathematics Studies, No. 61. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1968.
- [22] Katura Miyazaki. A note on genera of band sums that are fibered. *J. Knot Theory Ramifications*, 27(12):1871002, 3, 2018.
- [23] C. D. Papakyriakopoulos. On Dehn’s lemma and the asphericity of knots. *Ann. of Math. (2)*, 66:1–26, 1957.
- [24] Martin Scharlemann. Smooth spheres in $\mathbf{R}^4$ with four critical points are standard. *Invent. Math.*, 79(1):125–141, 1985.
- [25] Martin Scharlemann and Abigail Thompson. Link genus and the Conway moves. *Comment. Math. Helv.*, 64(4):527–535, 1989.
- [26] John Stallings. On topologically unknotted spheres. *Ann. of Math. (2)*, 77:490–503, 1963.
- [27] Shin’ichi Suzuki. Knotting problems of 2-spheres in 4-sphere. *Math. Sem. Notes Kobe Univ.*, 4(3):241–371, 1976.
- [28] E. C. Zeeman. Twisting spun knots. *Trans. Amer. Math. Soc.*, 115:471–495, 1965.